

# E-Z FLO BOTRAN® 6 DUST

**WARNING:**  
**KEEP OUT OF THE REACH OF CHILDREN**

635-485  
**ACCEPTED**  
Mar 5 1971  
FEDERAL INSECTICIDE  
AND FUNGICIDE ACT  
EPA REG. NO. 100-100-0101-101

**ACTIVE INGREDIENTS:**

2,6-Dichloro-4-nitroaniline ..... 6.0%

**INERT INGREDIENTS:** ..... 94.0%

100.0%

©Tuco Products Company — Division of the Upjohn Company  
U. S. Pat. No. 3,119,736

**DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.**

## DIRECTIONS FOR USE

### DIRECTIONS FOR USE ON EDIBLE CROPS

Crop	Disease	Dust Lb. Per Acre	Use
Peach and Nectarine	Blossom blight <i>Monilinia</i>	60	Apply at pink bud and full bloom.
	Fruit decay <i>Monilinia</i> Brown rot <i>Rhizopus</i> rot	60	Apply 18 days, 10 days and 1 day prior to harvest.
Sweet Cherry	Blossom blight <i>Monilinia</i>	50	Apply at popcorn, bloom, full bloom and petal fall.
	Fruit decay <i>Monilinia</i> Brown rot <i>Rhizopus</i> rot	50	Apply 10 days and 1 day prior to harvest.
Grape	<i>Botrytis</i> rot	30	Begin application while clusters are still loose and repeat application at two-week intervals. Allow one day between application and harvest.

### DIRECTIONS FOR USE ON EDIBLE CROPS — Continued

Crop	Disease	Dust Lb. Per Acre	Use
Tomato	Stem canker <i>Botrytis</i> (Greenhouse)	30	Dust stem of plant from ground level up to a height of 18 to 24 inches. Repeat weekly or as necessary.
Onion and Garlic	White rot <i>Sclerotium</i>	250-400	<b>Direct seeding</b> — Incorporate BOTRAN in top 1 1/2 inches of soil before onions are seeded. <b>Planting sets or buds</b> — Dust BOTRAN into furrow when planting so treated soil surrounds sets or buds. Rate/acre will vary in local areas.
SEE NOTE BELOW			
Lettuce (leaf, only)	Blight <i>Botrytis</i> rot (Greenhouse)	30	Apply dust 10 days following transplanting. Repeat applications when plants are half mature. Do not apply to wilted plants, seedlings or later than 14 days before harvest.
	(head)	Wilt <i>Botrytis</i> rot <i>Sclerotinia</i> (Field)	30

**NOTE:** Do not plant spinach as a follow-up crop in treated soil. These rates are not recommended for the North Central States.

### DIRECTIONS FOR USE ON ORNAMENTAL CROPS

Crop	Disease	Dust Lb. Per Acre	Use
Rose, Hydrangea	Blight <i>Botrytis</i>	30	<b>Field</b> —Apply BOTRAN prior to shipment or storage. <b>Stored</b> —Thoroughly dust storage area and all canes and dead foliage; repeat application during long storage periods.
Chrysanthe- mum, Geranium, Rose	Blight <i>Botrytis</i>	30	<b>Stock cuttings or greenhouse plants</b> — Begin when disease is anticipated or first appears. Dust foliage and flowers at 7- to 14-day intervals. During prolonged damp, cool periods, apply every 5 to 7 days.
Gladiolus	Dry rot <i>Stromatinia</i> (Soil)	250 (Broadcast)	Dust into furrow at planting time, so treated soil surrounds corms.
	Blight <i>Botrytis</i>	30	Dust thoroughly. Force fungicide between spikes. Apply to bunched cut flower spikes after harvest and before shipment or storage.

## WARNING

**Warning:** Keep this product out of the reach of children. Harmful if swallowed. Avoid inhalation of dust. Avoid contact with skin, eyes, or clothing. In case of contact, remove contaminated clothing and wash skin with soap and water. For eyes, flush with water. Avoid contamination of feed or foodstuffs.

**Precautions:** Do not use on germinating seeds or annual seedlings except as suggested.

This product is toxic to fish. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label. Do not apply when weather conditions favor drift of dust from crop areas treated. Apply this product only as specified on this label.

Net Contents and Manufacturer  
On Container