### WARNING:

## KEEP OUT OF THE REACH OF CHILDREN

#### **ACTIVE INGREDIENTS:**

| Parathion*         | 1.5%   |
|--------------------|--------|
| Sulphur            | 80.0%  |
| INERT INGREDIENTS: |        |
|                    | 100.0% |

\*O, O-Diethyl O-p-Nitrophenyl Thiophosphate.

### DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

# **DIRECTIONS FOR USE**

For the control of brown rot, scab, curculio and aphids on peaches apply at the rate of 25 to 40 pounds per acre. Two to six applications are usually necessary for curculio. Apply at pink, bloom and petal fall and every 10 to 14 days thereafter until control is achieved.

NOTE: Do not apply parathion to peaches closer than 14 days before harvest. Apply when the air is calm to avoid drift to other crops and premises within, surrounding or adjacent to the orchard.

**CAUTION:** No more than 5 lbs. actual parathion should be applied per acre per year to peaches. In California, do not apply more than once after bloom nor within 21 days of harvest.

For further information as to timing of applications consult nearest Federal or State Experimental Station.

**CONTAINER DISPOSAL:** Rinse equipment and containers and dispose of wastes and spillage by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning (keep out of smoke).

53-263-637

Settacturer

Container

ACCOPTED

WARNING: May Be Fatal if Swallowed, Inhaled or Absorbed Through Skin! Rapidly Absorbed Through Skin! Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

Not for use or storage in or around the home.

ANTIDOTE: ATROPINE IS AN ANTIDOTE — CONSULT PHYSICIAN FOR EMERGENCY SUPPLY.

FIRST AID TREATMENT: CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED PARATHION POISONING.

If symptoms or signs of poisoning include blurred vision, abdominal cramps and tightness in the chest, don't wait for a doctor but give two atropine tablets (each 1/100 gr.) at once. Remove patient immediately from the area where parathion is present. Remove contaminated clothing and wash the skin clean with plenty of soap and water to remove all traces of parathion. If swallowed, induce vomiting by giving warm salty or soapy water. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

PHYSICIAN'S NOTE: Warning symptoms include weakness, headache, tightness in the chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

**TREATMENT:** Give atropine, preferably by injection, grains 1/100 two or three tablets at once and parenterally or orally every hour as required up to 30 tablets or until pupils dilate. 2-PAM is also antidotal and may be administered in conjunction with atropine. Never give morphine. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood tests.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label. Birds feeding on treated areas may be killed.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

U.S.D.A. Reg. No. 635-491

ACCEPTED WITH COMMENT'S

09-18