

ACCEPTED

CONTAINER

# E-Z FLO METHYL PARATHION 1.5 THIODAN® 3 E.C.

(Contains 1.5 Lbs. Methyl Parathion and 3 Lbs. Thiodan\* Per Gallon)

**DANGER:**

KEEP OUT OF THE REACH OF CHILDREN

For Additional Warnings, See Appropriate State Laws



**POISON**



**ACTIVE INGREDIENTS:**

0,0-dimethyl 0-p-nitrophenyl thiophosphate 15.9%

Endosulfan\* 31.9%

Xylene 39.5%

**INERT INGREDIENTS:** 12.7%

100.0%

\*Hexachlorocyclohexane  
\*Thiodan is a registered trademark of Canadian Hoechst Limited  
U.S. Patent No. 2,799,685

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

SOLD BY

## E-Z FLO CHEMICAL COMPANY

DIVISION OF GROWER SERVICE CORPORATION

ABERDEEN, NORTH CAROLINA

COLUMBUS, OHIO

LANSING, MICHIGAN

53-263 B34

EPA Reg. No. 131-2

### DIRECTIONS FOR USE

Methyl Parathion 1.5 Thiodan 3 E.C. is designed for control of certain insects on cotton and tobacco. Use as directed below. If used in aerial application, use 10 gal. of water per acre.

**COTTON:** Use 1 1/2 pints in adequate water for coverage per acre for control of Cotton Leafworms and Thrips. Use 1 1/2 pints to 2 pints in adequate water for coverage per acre for control of Boll Weevils, Cotton Aphids, Cotton Fleahoppers, Lygus Bugs and other Mounds and Spider Mites (Carmine and Desert). Use 2 pints in adequate water for coverage per acre for control of Bollworms, Cabbage Loopers, Flea Armyworms, Cotton Leaf Perforator and Stink Bugs.

Workers entering fields within 24 hours after application should wear protective clothing. Do not apply after buds open nor within 5 days of handpicking. Do not graze dairy or meat animals in treated fields.

**TOBACCO:** For the control of Flea Beetles, Hornworms, Budworms, Stink Bugs, Green Peach Aphids and Cabbage Loopers in the field, use 1 1/2 to 2 1/2 pints per acre in adequate water for coverage. In the seed bed, use 1 1/2 pints in 100 gallons of water. Apply 6 gallons of finished spray per 100 square yards of tobacco seed bed. Do not apply within 5 days of transplanting, 5 days of pruning or 15 days of harvest of tobacco. For control of Green June Beetle larvae in the seed bed, use 1/2 pint in 100 gallons of water. Apply 1 gallon of the finished spray per square yard of plant bed at first sign of larval activity.

### General Information

Application of Methyl Parathion 1.5 Thiodan 3 E.C. should begin when insect infestation reaches the point where control is desired. Full coverage of plants during application is essential for good control. The material should be used for the insects listed above at the rate given in sufficient water to provide uniform coverage. Application should be made 3 to 5 day intervals depending upon the severity of the infestation and local conditions. To mix, fill the spray tank 1/4 to 1/2 full of water and with the agitator running, add the required amount of this material, then finish filling the spray tank to the desired level with water.

Application of this product to tobacco should be made only by a trained operator.

Rinse equipment and containers, and dispose of wastes by burying them in unproductive lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning (stay out of smoke).

**DESTROY CONTAINER WHEN EMPTY. NEVER RE USE**

# E-Z FLO METHYL PARATHION 1.5 THIODAN 3 E.C.

Contains 1.5 lbs. Methyl Parathion and 3 lbs. Thiodan Per Gallon

<b>ACTIVE INGREDIENTS</b>	
O,O dimethyl O-p-nitrophenyl thiophosphate	15.9%
Endosulfan*	31.9%
Xylene	39.5%
<b>INERT INGREDIENTS</b>	12.7%
	100.0%

REGISTERED TRADEMARKS OF THE MANUFACTURERS OF THE ACTIVE INGREDIENTS

SOLD BY  
**E-Z FLO CHEMICAL COMPANY**  
 DIVISION OF GROWER SERVICE CORPORATION  
 GREENSBORO, NORTH CAROLINA    COLUMBUS, OHIO    LANSING, MICHIGAN

## DIRECTIONS FOR USE

### APPLICATION

### TOBACCO

## General Information

DESTROY CONTAINER WHEN  
 EMPTY. NEVER RE-USE.