RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

STOP

METHYL PARATHION 1.5 THIODAN® 3 E.C.

DANGER:

KEEP OUT OF REACH OF CHILDREN

See Side Panel for additional Warnings

POISON

POISON

ACCEPTED

SEP 9 1978

Under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 635-55

SOLD BY

Lansing, Michigan 48901

DIRECTIONS FOR USE

Methyl Parathion 1.5 Thiodan 3 E.C. is designed for control of certain insects on cotton and tobacco when used as directed below. If used in aerial application use 10 gallons of water per acre.

COTTON: Use 1½ pints in adequate water for coverage per acre for control of Cotton Leafworms and Thrips. Use [illegible] pints to [illegible] pints in adequate water for coverage per acre for control of Boll Weevils, Cotton Aphids, Cotton Fleahoppers, Lygus Bugs and other Mirids and Spider Mites (Carmine and Desert). Use 2 pints in adequate water for coverage per acre for control of Bollworms, Cabbage Loopers, Fall Armyworms, Cotton Leaf Perforator and Stink Bugs.

Workers entering fields within [illegible] hours after application should wear protective clothing. Do not apply after [illegible] within [illegible] days of handpicking. Do not graze [illegible] animals in treated fields.

TOBACCO: For the control of Flea Beetles, [illegible], Stink Bugs, Green Peach Aphids and Cabbage Loopers [illegible] 2½ pints per acre in adequate water for coverage [illegible] use 1½ pints in 10 gallons of water [illegible] per 10 square yards of tobacco [illegible] [illegible] Green June Beetles [illegible] gallons of water. Apply [illegible] first sign [illegible]

DANGER

[illegible]

FIRST AID TREATMENT

ANTIDOTE [illegible]

[illegible]

General Information

Application of Methyl Parathion 1.5 Thiodan 3 E.C. should begin when insect infestation reaches the point where control is desired. Full coverage of plants during application is essential for good control. The material should be used for the insects listed above at the rate given in sufficient water to provide uniform coverage. Application should be made 3 to 5 day intervals depending upon the severity of the infestation and local conditions. To mix [illegible] the spray tank [illegible] full of water and [illegible] the agitator running [illegible] the required amount of this material [illegible] the spray [illegible] with water.

[illegible]

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.