

E-Z FLO 5% ROTENONE E. C.

(CONTAINS 0.4035 LBS. ROTENONE PER GALLON)

WARNING:

KEEP OUT OF THE REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL WARNINGS

ACTIVE INGREDIENTS:

Rotenone	5.00%
Other cube resins	6.12%
Methylated naphthalene	78.88%

INERT INGREDIENTS:

10.00%	
<hr/>	
100.00%	

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

SOLD BY

E-Z FLO CHEMICAL COMPANY

DIVISION OF GROWER SERVICE CORPORATION

ABERDEEN, NORTH CAROLINA COLUMBUS, OHIO LANSING, MICHIGAN

53-263-884

U.S.D.A. Reg. No. 635-515

DIRECTIONS FOR USE

CATTLE GRUBS (ALL CATTLE)

Dilute one gallon of this product with 100 gallons of water. Use approximately 2 quarts of the diluted spray per animal, spraying at 400 lb. pressure at a distance of 12-16 inches from backs of animals.

Low pressure hand sprayers may be used by thoroughly wetting backs of animals and scrubbing backs with brush to penetrate the cysts.

The first application should be made when cysts containing live grubs appear in the backs. Repeat at 30-day intervals. Two or three applications are usually sufficient.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

WARNING

Poisonous if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. In case of contact, wash immediately with soap and water. Wash all contaminated clothing with soap and hot water before re-use.

This product will kill fish. Keep out of lakes, ponds, and streams.

635-515

E-Z FLO 5% ROTENONE E. C.

(CONTAINS 0.4035 LBS. ROTENONE PER GALLON)

WARNING:

KEEP OUT OF THE REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL WARNINGS

ACTIVE INGREDIENTS:

Rotenone 5.00%

Other cube resins 6.12%

Methylated naphthalene 78.88%

INERT INGREDIENTS:

10.00%

100.00%

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

SOLD BY

E-Z FLO CHEMICAL COMPANY

DIVISION OF GROWER SERVICE CORPORATION

ABERDEEN, NORTH CAROLINA COLUMBUS, OHIO LANSING, MICHIGAN

53-263-884

U.S.D.A. Reg. No. 635-515

DIRECTIONS FOR USE

CATTLE GRUBS (ALL CATTLE)

Dilute one gallon of this product with 100 gallons of water. Use approximately 2 quarts of the diluted spray per animal, spraying at 400 lb. pressure at a distance of 12-16 inches from backs of animals.

Low pressure hand sprayers may be used by thoroughly wetting backs of animals and scrubbing backs with brush to penetrate the cysts.

The first application should be made when cysts containing live grubs appear in the backs. Repeat at 30-day intervals. Two or three applications are usually sufficient.

WARNING

Poisonous if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. In case of contact, wash immediately with soap and water. Wash all contaminated clothing with soap and hot water before re-use.

This product will kill fish. Keep out of lakes, ponds, and streams.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.