E-Z FLOEGO MOUSE BAIT

JUL : 1973

FUNCICIDE AND RESEARCH FOR ECONOMIC POISON REGISTER ED UNGER NO. 735 -522

Net Contains and Manufacturer On Container

WARNING:

KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENT:

Zinc Phosphide 2.0%
INERT INGREDIENTS: 98.0%
100.0%

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

An aid in the control of Meadow Mice, White Footed Mice; in orchards, nurseries, berry farms, and truck gardens.

BROADCAST BAITING. Can be applied by cyclone seeder, or by hand. By hand, throw tablespoon amounts into heavy cover along tree rows, rock outcrops, and fence lines. Apply at the rate of 6-10 pounds per acre. Do not broadcast over growing crops.

CONTAINER AND SPILLAGE DISPOSAL. In case of spillage, collect and bury in a non-crop area away from water supplies. Burn or bury empty bags (stay out of smoke).

WARNING

May be fatal if swallowed. Do not breathe dust or fumes. Do not contaminate feed and foodstuffs. Keep away from children and domestic animals. Avoid contact with skin. Wash hands after using. Wear rubber gloves when handling bait. Store in dry place.

ANTIDOTE

Give one tablespoon of salt in a glass of warm water; repeat until vomit fluid is clear. Have patient lie down and keep warm. Call a physician immediately.

SPECIAL CAUTIONS: Do not allow bait to be exposed on bare ground. Store poisoned grains in locations not accessible to children, pets, poultry and domestic animals. Do not apply to bare ground. To protect wildlife, scatter bait evenly or hide under cover in runways. Do not broadcast over growing crops.

This product is toxic to fish, birds, and other wildlife. Treated seeds exposed on soil surface may be hazardous to birds and other wildlife. Treated baits should be placed in locations not accessible to children, pets, wildlife and domestic animals, or in tamper-proof bait boxes. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.