

E-Z FLO TRITHION® 4-E

(Contains 4 Lbs. Trithion Per Gallon)

WARNING:

KEEP OUT OF THE REACH OF CHILDREN
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

ACTIVE INGREDIENTS:

Carbophenothion [S-[(p-Chlorophenylthio)methyl]

O,O-Diethyl Phosphorodithioate]

45.8%

Xylene

45.0%

INERT INGREDIENTS:

9.2%

100.0%

U. S. Pat. No. 2,793,224.
*T.M. Stauffer Chemical Company for insecticide-acaricide

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

SOLD BY

E-Z FLO CHEMICAL COMPANY

DIVISION OF GROWER SERVICE CORPORATION

ABERDEEN, NORTH CAROLINA

COLUMBUS, OHIO

LANSING, MICHIGAN

06-18

53-263-895

U.S.D.A. Reg. No. 635-525

DIRECTIONS FOR USE

Application should be made only by experienced personnel using approved safety equipment.

Pour recommended amount of this material into nearly filled spray tank, add balance of water to fill tank. Keep agitator running during filling, mixing, and spraying operations. Do not allow mixture to stand. Do not combine emulsifiable liquids and wettable powders in same spray tank unless previous use of the materials being combined has proved them to be physically compatible.

RECOMMENDATIONS FOR DORMANT SPRAYS

Unless otherwise specified, rates in this section are given in terms of Trithion 4-E and spray oil per 100 gals. of water for use in a thorough coverage spray. If concentrate sprays are used, the amount of Trithion 4-E per 100 gals. should be adjusted in order to apply the same amount of Trithion per acre. Apply a minimum of 1 quart Trithion 4-E per acre.

APPLES, CRABAPPLE, PEAR, QUINCE: Overwintering aphid and mite eggs, pear psylla, scales (Forbes, oyster shell, Putnam, San Jose, Scurfy) — Use ½ - 1 pint plus 1 - 2 gals. spray oil with thorough coverage. Do not apply more than 1½ qts. Trithion 4-E per acre to crabapple, pears, or quinces nor more than 4½ qts. Trithion 4-E to apples.

APRICOTS: Overwintering mite eggs, scales (European fruit lecanium, San Jose, soft brown) — Use ½ - 1 pint plus 1 - 2 gals. spray oil. Do not apply more than 2 qts. Trithion 4-E per acre.

CHERRY: Overwintering aphid and mite eggs, scales (European fruit lecanium, Forbes, olive (Parlatoria) San Jose) — Use ½ - 1 pint plus 1 - 2 gals. spray oil. Do not apply more than 2 qts. Trithion 4-E per acre.

PEACH, NECTARINE: Scales (European fruit lecanium, olive (Parlatoria), San Jose) — Use ½ - 1 pint plus 1 - 2 gals. spray oil. Do not apply more than 2 qts. Trithion 4-E per acre.

PLUM, PRUNE: Overwintering aphid and mite eggs, scales (European Fruit lecanium, olive (Parlatoria) San Jose) — Use ½ - 1 pint plus 1 - 2 gals. spray oil. Do not apply more than 2 qts. Trithion 4-E per acre.

WALNUT: Walnut aphid, overwintering mite eggs, scale (calico, European fruit lecanium, frosted) — Use 1 - 2 quarts plus 2 gals. spray oil per acre in sufficient water for coverage. Do not apply more than 2 gals. of oil nor 2 qts. of Trithion 4-E per acre.

RECOMMENDATIONS FOR VEGETABLE, VINE AND FIELD CROPS

Dosage rates are given in terms of TRITHION 4-E per acre in sufficient water to provide good coverage.

BEANS (Succulent) Snap, Lima: Bean aphid, Mexican bean beetle, potato leafhopper, serpentine leaf miner, two-spotted mites — Use 1 ½ pints. Do not apply within 7 days of harvest. Do not graze or feed crop residue to livestock.

COTTON (For Cotton South): Early Season: Carnine (cinnabar), mite, two spotted mite, tropical mite — Use ½ - 1 pint per acre in sufficient water for thorough coverage.

WARNING:

DO NOT GET IN EYES, ON SKIN, OR ON CLOTHING. IN CASE OF CONTACT, WASH IMMEDIATELY WITH SOAP AND WATER. FOR EYES FLUSH WITH WATER FOR 15 MINUTES. WASH HANDS, ARMS AND FACE THOROUGHLY WITH SOAP AND WATER, BEFORE EATING OR SMOKING. WASH CONTAMINATED CLOTHING WITH SOAP AND HOT WATER BEFORE RE-USE.

TRITHION upon repeated, prolonged and careless use, may cause cholinesterase inhibition. ATROPINE IS AN ANTIDOTAL. Commercial operators should keep supply of Atropine 1:100 grain tablets. 2-PAM is also antidotal and may be administered in conjunction with atropine.

FOR COMMERCIAL APPLICATION, WEAR RUBBER GLOVES, PROTECTIVE CLOTHING, RUBBER OVERSHOES AND GOGGLES.

OBSERVE THE FOLLOWING USE PRECAUTIONS

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

DO NOT APPLY OR ALLOW TO DRIFT TO AREAS OCCUPIED BY UNPROTECTED HUMANS, BENEFICIAL ANIMALS OR ONTO ADJOINING FOOD AND FORAGE CROPS.

THIS PRODUCT IS TOXIC TO FISH AND WILDLIFE. KEEP OUT OF LAKES, STREAMS AND PONDS. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

THIS PRODUCT IS TOXIC TO BEES AND SHOULD NOT BE APPLIED WHEN BEES ARE ACTIVELY VISITING THE AREA.

THE FEDERAL FOOD AND DRUG ADMINISTRATION HAS SET OFFICIAL TOLERANCES FOR CERTAIN PESTICIDE RESIDUES. IN ORDER THAT RESIDUES ON FOOD AND FORAGE CROPS WILL NOT EXCEED ESTABLISHED TOLERANCES, USE ONLY AT RECOMMENDED RATES, AND DO NOT APPLY CLOSER TO HARVEST THAN SPECIFIED.

DIRECTIONS (Continued)

Mid and Late Season: ¾ - 1 pint per acre in sufficient water for thorough coverage. If mite populations are high use 1 - 1½ pints per acre in sufficient water for thorough coverage and repeat application in 5 - 10 days.

Cotton leafworm — Use ½ - 1 pint per acre in sufficient water for thorough coverage.

Cotton leaf perforator, fleahopper, aphids — Use 1 qt. per acre in sufficient water for thorough coverage.

NOTE: Do not graze dairy or meat animals in treated cotton fields.

FOREST AND CHRISTMAS TREE PLANTINGS: European pine shoot moth and tip moth. Apply 1½ - 2 pints in 5 - 50 gals. of water per acre. Repeat application if necessary.

LAWNS, ORNAMENTAL TURF: Chinch bugs — Wet lawn thoroughly, then apply 1½ pints in 100 - 150 gal. water to 5,000 sq. feet of lawn area. Do not permit children or domestic animals on lawn until area is completely dry. Birds feeding on treated areas may be killed.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

9-19-70
635-525

NET CONTENTS ON CONTAINER

