

E-Z FLO COPPER, SEVIN*, SULPHUR 3.4-5-70 DUST

CAUTION:
KEEP OUT OF THE REACH OF CHILDREN

ACCEPTED
 10-22-70
 635-548

ACTIVE INGREDIENTS:

Sulphur (95% through 325 Mesh Screen).....	70.0%
Carbaryl (1-naphthyl N-methylcarbamate)	5.0%
**Copper as Metallic	3.4%
INERT INGREDIENTS	21.6%
	100.0%

*Trade Mark of UCC for 1-naphthyl N-methylcarbamate. U. S. Patent Nos. 2,903,478; 3,009,855
 **Copper is in form of Cuprous Oxide.

Net Contents and Manufacturer
 On Container

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE ON PEANUTS

For the control of leaf spot on peanuts apply at 7 to 10 day intervals beginning when spots first appear on basal leaves at the rate of 15 to 20 pounds per acre.

ARMYWORMS: Velvet Bean Caterpillar (or peanut worms) on peanuts: begin applications at the rate of 20 to 30 pounds per acre when small worms first appear.

THRIPS on peanuts: Begin applications at the rate of 20 to 30 pounds per acre when small thrips first appear. Repeat the applications in 7 to 10 days. Apply additional dustings during the season as needed.

CAUTION

Harmful if swallowed or inhaled. Avoid breathing dust. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.