

E-Z FLO CYPREX[®] PARATHION 3-3 DUST

DANGER.

KEEP OUT OF THE REACH OF CHILDREN



POISON



USE PRECAUTIONS

Do not prune fruit trees or thin fruit within 15 days after application. Do not apply during bloom when honey bees or other pollinators are active. Avoid contact with dust drift on flowers, shrubs, or fruits, vegetables, and forage crops. Keep children and animals off treated areas until chemical has been washed into the soil by sprinkling or rain and the grass is dry.

Do not re-use container. Destroy by burying in a safe place. Do not burn.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates and do not apply closer to harvest than specified. Consult your State Agricultural Experiment Station for specific information consistent with labeling as the timing, number and rate of applications may vary with local conditions.

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

ACTIVE INGREDIENTS:

Dodine (n-Dodecylguanidine acetate) 3.0%

Parathion (O,O-Diethyl O-p-Nitrophenyl Thiophosphate) 3.0%

INERT INGREDIENTS: 94.0%

100.0%

*Cyprex is a registered trademark of American Cyanamid Company U. S. Patent No. 2,867,567

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

APPLES: For the control of Scab, Codling Moth, Plum Curculio, Orange Tortrix, San Jose Scale, Forbes Scale, Scurfy Scale, Red Banded Leaf Roller, Apple Maggot, Woolly Apple Aphid, Bud Moth, Fruit Tree Leaf Roller, Rosy Apple Aphid, Green Apple Aphid, Leafhopper, European Red Mite and Two-spotted Mite. Apply 35-45 lbs. per acre at 7 day intervals or as needed to maintain control from pre-bloom period through 1st cover. Repeat at 7-10 day intervals throughout the summer months. Make full coverage applications. Do not apply within 14 days of harvest. Do not apply to susceptible varieties such as McIntosh, Cortland and related varieties, Golden Delicious, Grimes Golden, Rhode Island Greening or Jonathan.

CHERRIES: For the control of Leaf Spot, Aphids, Mites, Leafminer, Plum Curculio, Cherry Fruit Worm, Oriental Fruit Moth, Bud Moth, Tortrix, Cankerworm, San Jose Scale. Apply 35-45 lbs. per acre beginning at petal fall or 5 days after first fruit fly emerges. Repeat every 7-10 days and immediately after every rain. Make full coverage application. Do not apply within 14 days of harvest. Make a post-harvest application for additional protection.

American Cyanamid Company
 260 North Zeeb Road
 York, Pennsylvania 17403

WARNING

Extremely poisonous if swallowed. Irritated on contact through the eye. Avoid getting dust on the skin or the eyes or on clothing. Rapidly absorbed through skin. Do not breathe dust. Wear a respirator which has been tested by the U. S. Department of Agriculture for parathion protection. Avoid contact with clothing, footwear, rubber gloves, hat, and other items which may be contaminated by this product. Wash hands, skin, and hair with soap and water thoroughly after application or before eating or drinking. Wash contaminated clothing with hot water and soap before reuse. Do not use contaminated equipment. Do not use in the operator's home or in areas where children may be present. Do not use in areas where children may be present. Do not use in areas where children may be present.

FIRST AID TREATMENT

IF SWALLOWED: Do not induce vomiting. Give 1-2 glasses of water. If unconscious, give water by mouth.

INTERNAL USE

IF SWALLOWED: Do not induce vomiting. Give 1-2 glasses of water. If unconscious, give water by mouth.

EYES

IF IRRITATED: Flush eyes with water for 15 minutes.

SKIN

IF IRRITATED: Wash with soap and water.

ACCEPTED

E-Z FLO CYPREX, PARATHION 3-3 DUST



USE PRECAUTIONS

Do not prune fruit trees or thin fruit within 15 days after application. Do not apply during bloom when honey bees or other pollinators are active. Avoid contact with dust drift on flowers, shrubs, or fruits, vegetables, and forage crops. Keep children and animals off treated areas until chemical has been washed into the soil by sprinkling or rain and the grass is dry.

Do not reuse container. Destroy by burying in a safe place. Do not burn.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates and do not apply closer to harvest than specified. For a list of State Agricultural Experiment Stations for specific information, contact with laboratory, the timing, number and rate of applications, and any other local conditions.

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Do not contaminate any body of water or apply when conditions favor drift from the crop area treated.

This product is highly toxic to bees and other insects that are beneficial on crops. Protective information should be obtained from your State Department of Agricultural Extension Service.

WARNINGS

ACTIVE INGREDIENTS:

Dodine (n-Dodecylguanidine acetate)	3.0%
Parathion (O,O-Diethyl O-p-Nitrophenyl Thiophosphate)	3.0%

INERT INGREDIENTS:

94.0%
100.0%

E-Z Flo Cyprex is a registered trademark of American Cyanamid Company, U.S. Patent No. 2,747,556.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

Apply to all susceptible fruit trees, including Malus, Pyrus, Prunus, and Pisonia, and to all susceptible shrubs, including Rubus, and to all susceptible forage crops, including alfalfa, clover, and lucerne. Apply to all susceptible vegetables, including beans, peas, and tomatoes. Apply to all susceptible ornamental plants, including roses, geraniums, and petunias. Apply to all susceptible weeds, including quackgrass and wild oats. Apply to all susceptible insects, including aphids, beet beetles, and sawflies. Apply to all susceptible diseases, including apple scab and cherry leaf spot.

CHERRIES

Apply to all susceptible cherry trees, including Prunus, and to all susceptible shrubs, including Rubus. Apply to all susceptible forage crops, including alfalfa, clover, and lucerne. Apply to all susceptible vegetables, including beans, peas, and tomatoes. Apply to all susceptible ornamental plants, including roses, geraniums, and petunias. Apply to all susceptible weeds, including quackgrass and wild oats. Apply to all susceptible insects, including aphids, beet beetles, and sawflies. Apply to all susceptible diseases, including cherry leaf spot.