

DANGER:
KEEP OUT OF THE REACH OF CHILDREN



POISON



ACTIVE INGREDIENTS:

Dodine (n-Dodecylguanidine acetate)	3.0%
Parathion (O,O-Diethyl O-p-nitrophenyl thiophosphate)	3.0%
Lead Arsenate (PbHAsO₄)*	15.0%

INERT INGREDIENTS:	79.0%
	100.0%

* Cyprex is a registered trademark of American Cyanamid Company. U. S. Patent No. 2,867,562.

* Total arsenic expressed as metallic 21.14%. Arsenic in water soluble form, expressed as metallic, not more than 0.35%. Lead expressed as metallic 58.00%.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

APPLES: For the control of Scab, Codling Moth, Plum Curculio, Orange Tortrix, San Jose Scale, Forbes Scale, Scurfy Scale, Red Banded Leaf Roller, Apple Maggot, Woolly Apple Aphid, Bud Moth, Fruit Tree Leaf Roller, Rosy Apple Aphid, Green Apple Aphid, Leafhopper, European Red Mite and Two spotted Mite. Apply 35-45 lbs per acre at 7 day intervals or as needed to maintain control from pre-bloom period through 1st cover. Repeat at 7-10 day intervals throughout the summer months. Make full coverage applications. Do not apply within 30 days of harvest. Do not use treated apples in the manufacture of apple pomace for use in livestock feeds. Remove excess residue at harvest. Do not apply to susceptible varieties such as McIntosh, Cortland and related varieties, Golden Delicious, Grimes Golden, Rhode Island Greening or Jonathan.

CHERRIES: For the control of Leaf Spot, Aphids, Mites, Leafminer, Plum Curculio, Cherry Fruit Worm, Oriental Fruit Moth, Bud Moth, Tortrix, Cankerworm, San Jose

Scale. Apply 35-45 lbs per acre beginning at petal fall or 5 days after first fruit fly emerges. Repeat every 7-10 days and immediately after every rain. Make full coverage application. Do not apply within 14 days of harvest on cherries to be processed, nor 30 days on cherries for fresh market. Remove excess residue at harvest. Do not graze treated forage by livestock. Make a post harvest application for additional protection.

USE PRECAUTIONS

Do not prune fruit trees or thin fruit within 15 days after application. Do not apply during bloom when honey bees or other pollinators are active. Avoid contact with spray or dust drift on flowers, shrubs, or fruits, vegetables, and forage crops. Keep children and animals off treated areas until chemical has been washed into the soil by sprinkling or rain and the grass is dry.

Do not re-use container. Destroy by burying in a safe place. Do not burn.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates and do not apply closer to harvest than specified. Do not graze livestock on treated areas. Do not feed treated forage to livestock. Consult your State Agricultural Experiment Station for specific information consistent with labeling as the timing, number and rate of applications may vary with local conditions.

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated.

WARNING

FIRST AID TREATMENT:

INTERNAL

IF IN

EYES

INHALATION

Feb. 6, 1968 - Net contents and manufacturer on container.

POISON



ACTIVE INGREDIENTS:

Dodine (n-Dodecylguanidine acetate)	3.0%
Parathion (O,O-Diethyl O-p-nitrophenyl thiophosphate)	3.0%
Lead Arsenate (PbHAsO₄)*	15.0%

INERT INGREDIENTS:

79.0%
100.0%

Cyprex is a registered trademark of American Cyanamid Company, U.S. Patent No. 2,667,562.

*Total arsenic expressed as meta-arsenic acid. Arsenic in water soluble form expressed as meta-arsenic acid not more than 0.35%. Lead expressed as metallic lead.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

APPLES: For the control of Saprotrophic Rot, Fire Blight, and other diseases of the fruit and foliage of the apple tree. Apply to the foliage and fruit of the tree at the following rates: For Saprotrophic Rot, apply 1.5 to 2.0 lb. per acre; for Fire Blight, apply 1.5 to 2.0 lb. per acre; for other diseases, apply 1.5 to 2.0 lb. per acre. Apply to the foliage and fruit of the tree at the following rates: For Saprotrophic Rot, apply 1.5 to 2.0 lb. per acre; for Fire Blight, apply 1.5 to 2.0 lb. per acre; for other diseases, apply 1.5 to 2.0 lb. per acre.

CHERRIES: For the control of Saprotrophic Rot, Fire Blight, and other diseases of the fruit and foliage of the cherry tree. Apply to the foliage and fruit of the tree at the following rates: For Saprotrophic Rot, apply 1.5 to 2.0 lb. per acre; for Fire Blight, apply 1.5 to 2.0 lb. per acre; for other diseases, apply 1.5 to 2.0 lb. per acre.

USE PRECAUTIONS

Read the entire label carefully before using this product. Do not use if the label is torn, mutilated, or missing important instructions or warnings. Do not use if the product is past its expiration date. Do not use if the product is not in its original container. Do not use if the product is not in its original container.

Do not use if the product is past its expiration date. Do not use if the product is not in its original container. Do not use if the product is not in its original container. Do not use if the product is not in its original container. Do not use if the product is not in its original container. Do not use if the product is not in its original container.

Feb. 6, 1968 - Not contains and manufacturer on container.