

E-Z FLO BULK METHYL BROMIDE

DANGER
KEEP OUT OF REACH OF CHILDREN



POISON



ACTIVE INGREDIENTS

Methyl Bromide 68.6%

Chloropicrin 1.4%

INERT INGREDIENTS

10.0

100.0%

The manufacturer makes no warranties, express or implied, that the product will perform as described or that the results will extend beyond the description on the label. Use only as directed.

SOLD BY

E-Z FLO CHEMICAL COMPANY

DIVISION OF GROWER SERVICE CORPORATION

ABERDEEN, NORTH CAROLINA

COLUMBUS, OHIO

LANSING, MICHIGAN

LEXINGTON, KENTUCKY

53-263-907

A-111-N-635-077

DIRECTIONS FOR USE

PRODUCT USES

For pre-planting treatment of TOBACCO seed and plant beds, also GROUND FLORAL and NURSERY CROPS.

DIRECTIONS: Use Methyl Bromide for control of nematodes, soil-borne cyst formers, meadow nematode, root knot, tapeworm, sting, and spines. Also for the control of insects such as white grubs, wire worms, and other similar soil-borne pests. Apply when the soil is in a friable, moist condition.

APPLICATION: Use Methyl Bromide at the rate of 10 to 20 pounds per acre. Apply by means of nozzles spaced not more than 40 inches apart at a depth of 4 to 6 inches. Use at the recommended rate only on soils with high moisture content or after the temperature has risen to 50°F with protective cover. Do not apply to wet soil. Do not apply to soil with protective cover until the soil is dry. Do not apply to soil with protective cover until the soil is dry. Do not apply to soil with protective cover until the soil is dry.

PRECAUTIONS: Avoid breathing dust or fumes. Do not get in eyes. Avoid contact with skin. Wash thoroughly after handling. Do not eat, drink, or smoke while using. Do not use in confined spaces. Do not use in areas where there is a fire hazard. Do not use in areas where there is a risk of explosion.

PERSONAL PROTECTION: Wear protective clothing and shoes. Avoid contact with skin and eyes. Wash thoroughly after handling.

CONTAINER DISPOSAL

Empty containers should be returned to the manufacturer for disposal. Do not reuse.

DIRECTIONS (Continued)

Apply fertilizers containing nitrates or ammonium sulfate after the crop is established and the soil temperatures are above 65°F to avoid ammonia injury or nitrogen starvation.

To prevent injury to desirable living plants, keep edge of cover at least 18 inches from drip line. Do not apply to areas containing living roots of these plants.

Fumigated areas may be protected against recontamination by wind-er frames or trenches to prevent flooding or washing. Recontamination by footwear or equipment should be kept to a minimum.