

MADE IN U.S.A. (Contains 1 lb. Phosdrin and 2 lbs. Thiodan per gallon)

**POISON**

**DANGER: KEEP OUT OF THE REACH OF CHILDREN**


See Side Panel for Additional Warnings and Antidote Statements

**ACTIVE INGREDIENTS:**

Alpha isomer of 2-carbomethoxy-1-methylvinyl dimethyl phosphate ..... 6.94%  
 and related compounds ..... 4.02%  
 Endosulfan\* ..... 23.26%  
 Xylene ..... 60.72%

**INERT INGREDIENTS:** ..... 4.45%

**Total 100.00%**



\*Phosdrin is a registered trademark of Shell Chemical Company.  
 †Thiodan is a registered trademark of Canadian Hoechst Limited.  
 U. S. Patent No. 2,759,685.  
 \*Hexachlorocyclohexane-2, 4, 4-trichloro-1-hydroxy-1-oxide.

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

**SOLD BY**  
**E-Z FLO CHEMICAL COMPANY**  
 DIVISION OF KINGSO COMPANY  
 COLUMBUS, OHIO LANSING, MICHIGAN

5 50-033-280 U.S.D.A. Reg. No. 635072

NET CONTENTS ON CONTAINER

**DIRECTIONS FOR USE**

CROP	TO CONTROL	REMARKS
Cabbage Cauliflower Broccoli Cabbage Sprouts	Aphids, Lygus Bug, Loopers, Grasshoppers, Leafhoppers, Spittlebug Caterpillar, Cutworms (climber), Dipterous Leafminer (adults), Imported Cabbageworm, Stink Bugs, Diamond-back Moth Larvae, Cross-striped Cabbageworm, Harlequin Bug, Flea Beetle	Use 1 to 2 quarts in sufficient amount of water to cover one acre for all insects except Flea Beetles, Diamond-back Moth Larvae, Cross-striped Cabbageworm, Harlequin Bug and Flea Beetles. Use 1 1/2 to 2 quarts by spot application when insects first appear and repeat as required. Do not apply to cabbage or cauliflower within 7 days of harvest. Do not apply to cauliflower and Brassica sprouts within 10 days of harvest.
Cucumbers Squash (Squash, etc.)	Aphids, Flea Beetle, Grasshoppers, Leafhoppers, Mites	Use 1 to 2 quarts in sufficient amount of water to cover one acre for all insects except aphids. Use 1 quart per acre for spot applications when pests first appear and repeat as required. Do not apply within 1 day of harvest.
Emulsifol	Aphids, Flea Beetle, Colorado Potato Beetle, Whitefly, Green Stink Bug	Use 1 quart in sufficient amount of water to cover one acre. Apply when pests first appear and repeat as required. Do not apply within one day of harvest.
Miscellaneous Cauliflowers and Wintergreens	Aphids, Lygus Bug, Flea Beetle, Cutworm, Grasshoppers, Spittlebug Caterpillar, Cutworm (climber), Dipterous Leafminer (adults), Mites, Green and Red Flea	Use 1 to 2 quarts in sufficient amount of water to cover one acre for all insects except aphids. Use 1 1/2 to 2 quarts by spot application when pests first appear and repeat as required. Do not apply within one day of harvest.
Peas	Aphids, Cutworm, Leafhoppers, Mites	Use 1 quart in sufficient amount of water to cover one acre. Apply when pests first appear and repeat as required. Do not apply within one day of harvest.
Peas (Soybeans)	Aphids, Flea Beetle, Colorado Potato Beetle, Flea Beetle, Green Stink Bug, Imported Cabbageworm, Stink Bug, Flea Beetle	Use 1 to 2 quarts in sufficient amount of water to cover one acre for all insects except aphids. Do not apply to peas within 7 days of harvest. Do not apply to soybeans within 10 days of harvest.
Peas (Soybeans)	Aphids, Flea Beetle, Colorado Potato Beetle, Flea Beetle, Green Stink Bug, Imported Cabbageworm, Stink Bug, Flea Beetle	Use 1 to 2 quarts in sufficient amount of water to cover one acre for all insects except aphids. Do not apply to peas within 7 days of harvest. Do not apply to soybeans within 10 days of harvest.

**WARNINGS**

Prevention of death: In case of emergency, call your local poison center or hospital. If you are unable to do so, call the National Poison Center at Washington, D.C. (202) 638-2742. For more information, see the back of this container. Wash thoroughly with soap and water. For eyes, flush with plenty of water. Get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

**ANTIDOTE:** If swallowed: Give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. CALL A PHYSICIAN IMMEDIATELY.  
 If on skin: Remove contaminated clothing and wash skin immediately with soap and water. CALL A PHYSICIAN IMMEDIATELY.

**NOTE TO PHYSICIAN**

Phosdrin is cholinesterase inhibitor and treatment of severe poisoning should include the use of atropine; Thiodan produces central nervous system excitation which responds to the barbituric acid derivatives.  
 Enough atropine must be given in the shortest practical time to produce dilation of pupils. Patient must be under constant supervision and additional atropine may be necessary without delay. Clear chest by postural drainage. Use artificial respiration and oxygen if patient becomes comatose. Use of atropine may be necessary.

**USAGE CAUTIONS**

Keep on hand an emergency supply of 20 tablets, grains 1/200. In case of poisoning take 3 tablets (grs. 1/100) and call physician at once.  
 Keep all unprotected persons and animals out of operating areas or vicinity where there may be danger of drift. Vaccinated areas should not be re-entered on day of treatment. Do not store near feed and food products. Do not use in unventilated form. Keep containers closed. Leaking packages should be removed to a safe place. In case of spillage, wash down with large amounts of water. Decant both areas and equipment by washing with strong lye solution.  
**FLAMMABLE:** Keep away from heat and open flame. Do not store above 50°F or below 20°F.  
 This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Apply this product only as specified on this label.  
**DISPOSAL WARNINGS:** Do not reuse the container. Completely empty the contents and bury the residue chemical at least 18 inches deep in an isolated location away from water bodies. Break the container and bury the pieces or melt it in water deep in a water body away from water supply.  
 This product is highly toxic to bees. Do not apply to plants in bloom or treatment or residues on crops. For more information may be obtained from your Cooperative Agricultural Extension Service.