

E-Z FLO LEAD ARSENATE

DANGER:
KEEP OUT OF THE REACH OF CHILDREN



POISON



*Net Contents and Manufacturer
 On Container*

ACTIVE INGREDIENT: Lead Arsenate	96.0%
INERT INGREDIENTS:	4.0%
TOTAL	100.0%

Total arsenic, as metallic, not less than 20.7%; water soluble arsenic expressed as metallic, not more than 0.33%; lead as metallic 57%.

DIRECTIONS

FRUIT: On apples, pear, peach and plum to control codling moth, curculio, canker-worm and red-banded leaf roller, use at the rate of 2 pounds per 100 gallons of water. On tender foliage such as peach and plum add 8 pounds of Chemically Hydrated Lime to each 100 gallons of spray as a safener. Apply applications at petal fall and shuck-off and at 10 to 14 day intervals as necessary for control. Do not apply to any of the above fruit crops within 30 days of harvest. Remove all residues at harvest by washing, brushing or other effective means. Do not graze livestock in treated orchards.

635-594

VEGETABLES: On tomato to control hornworm and tomato fruit worm, spray using 4 pounds per 100 gallons of water. Do not exceed 250 gallons of spray per acre. Repeat as necessary for control. Remove all residue of material from edible parts of vegetables at harvest by thoroughly washing.

ORNAMENTALS: For control of bag worms on evergreens such as arborvitae and cedars, use 6 pounds per 100 gallons of water when insects are not over half grown. For larger insects use 8 pounds per 100 gallons of water. To control certain other chewing insects such as catalpa sphinx, walnut caterpillar, fall webworm, eastern tent caterpillar and forest tent caterpillar on shade trees, use at the rate of 4 pounds per 100 gallons of water.

DISPOSAL WARNING

Dispose of waste pesticide and container by burying in non-crop land away from water supplies. Never re-use container.



POISON



ANTIDOTE:

Give a tablespoon of salt in a glass of warm water and repeat until vomit fluids are clear. Then give 2 tablespoons of Epsom Salts or Milk of Magnesia in water and plenty of fresh water. Have victim lie down and keep quiet. Call a Physician immediately.

DANGER!

KEEP OUT OF REACH OF CHILDREN.

Poisons if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after using. Avoid breathing dust or spray mist. Keep away from pets, livestock, feed and foodstuffs.

PROTECT WILDLIFE: This product is toxic to wildlife. Keep out of lakes, streams or ponds. Do not apply where run off is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

PROTECT BEES: This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information is available from your Cooperative Agricultural Extension Service.

Do not use or store around the home.

E-Z FLO LEAD ARSENATE

DANGER:
KEEP OUT OF THE REACH OF CHILDREN



POISON



ACTIVE INGREDIENT: Lead Arsenate	95.0%
INERT INGREDIENTS:	4.0%
	TOTAL 100.0%

Total arsenic, as contained, not less than 30.7%, water soluble arsenic expressed as material, not more than 0.33%; lead as material 57%.

DIRECTIONS

FRUIT: On apples, pear, peach and plum to control codling moth, curculio, winter-worm and redbanded leaf roller, use at the rate of 2 pounds per 100 gallons of water. On tender foliage such as peach and plum add 8 pounds of Chemically Hydrated Lime to each 100 gallons of spray as a buffer. Apply applications at petal fall and shock-blossom at 10 to 14 day intervals as necessary for control. Do not apply to any of the above fruit crops within 30 days of harvest. Pick and market fruit at harvest by washing with clean water. This is an effective means. Do not pick untreated or treated orchards.

635-594

VEGETABLES: On tomato to control horn worm and tomato fruit worm, spray using 4 pounds per 100 gallons of water. Do not exceed 250 gallons of spray per acre. Repeat as necessary for control. Remove all residue of material from edible parts of vegetables at harvest by thoroughly washing.

ORNAMENTALS: For control of bag worms on evergreens such as arborvitae and cedars, use 6 pounds per 100 gallons of water when insects are not over half grown. For larger insects use 8 pounds per 100 gallons of water. To control certain other chewing insects such as catalpa sphinx, walnut caterpillar, fall webworm, eastern tent caterpillar and forest tent caterpillar on shade trees, use at the rate of 4 pounds per 100 gallons of water.

DISPOSAL WARNING

Dispose of waste pesticide and container by burying in non-crop land away from water supplies. Never re-use container.



POISON



ANTIDOTE;

DANGER!

Wash thoroughly with soap and water. If swallowed, induce vomiting. If inhaled, get fresh air. If on skin, wash with soap and water.

PROTECT WILDLIFE: This pesticide is extremely toxic to birds, fish, and aquatic life. Do not apply to water bodies or near water. Do not apply to areas where birds, fish, or aquatic life are present.

PROTECT BEES: Do not apply to flowering plants or crops when bees are active. Do not apply to areas where bees are present.