



E-Z Flo 57% Malathion E. C. (Premium Grade)

(Contains 5 Pounds Actual Malathion Per Gallon)

**CAUTION:
KEEP OUT OF REACH OF CHILDREN**

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

ACTIVE INGREDIENTS:

Malathion*	57.0%
Xylene	32.0%

INERT INGREDIENTS:	<u>11.0%</u>
	100.0%

* Dimethyl D-thiophosphate of Diethyl Mercaptosuccinate

The manufacturer makes no warranties, express or implied concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

SOLD BY

E-Z FLO CHEMICAL COMPANY

DIVISION OF GROWER SERVICE CORPORATION	
ABERDEEN, NORTH CAROLINA	LANSING, MICHIGAN
COLUMBUS, OHIO	LEXINGTON, KENTUCKY

CAUTIONS

Harmful if swallowed, avoid prolonged breathing of spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

ANTIDOTE: External: Wash with soap and water. **Internal:** Emetic of Mustard. If poisoning occurs, CALL A PHYSICIAN IMMEDIATELY. Artificial respiration and large doses of Atropine may be required.

This product is toxic to fish. Keep out of any body of water.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning (keep out of smoke).

SOLD BY

E-Z FLO CHEMICAL COMPANY

DIVISION OF GROWER SERVICE CORPORATION

ABERDEEN, NORTH CAROLINA
COLUMBUS, OHIO

LANSING, MICHIGAN
LEXINGTON, KENTUCKY

87-17

53-263-907

U.S.D.A. Reg. No. 635-605

out of any body of water.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning (keep out of smoke).

DIRECTIONS FOR USE

For protection of stored peanuts against infestations of red flour beetle, Indian meal moth, confused flour beetle, rice weevil, flat grain beetle, rusty grain beetle, lesser grain borer, granary weevil, and saw-toothed grain beetle, apply E-Z Flo 57% Malathion E. C. (Premium Grade) as follows:

Residual Warehouse Spray — Before Storing Peanuts: Clean warehouse thoroughly of trash and old remains of peanuts 1 to 2 weeks before new peanut crop is stored. Thoroughly spray with sufficient pressure interior of empty warehouse (including cracks and protected places) outside walls to height of 6 to 8 feet, and the ground to a distance of about 6 feet from warehouse by diluting 1 pint of E-Z Flo 57% Malathion E. C. (Premium Grade) in sufficient water to make 2½ gallons of spray or 1 gallon with 19 gallons of water. Apply finished spray at the rate of 2 gallons per 1,000 square feet or to run-off.

Bulk Spray Treatment — Peanuts Going Into Storage: Apply 2½ pints of E-Z Flo 57% Malathion E. C. (Premium Grade) in 5 gallons of water per 15 tons of farmers' stock peanuts as they go into storage. Apply coarse spray uniformly. Preferably use a suitable mechanical spray applicator which regulates rate of application to flow of peanuts. Otherwise, adjust pressure and size of nozzle opening to rate of flow of peanuts.

Supplemental Surface Spray: After warehouse has been filled, level surface of peanuts before spraying. Apply surface spray using 1¾ lbs. of E-Z Flo

DIRECTIONS (Continued)

Malathion WP (Premium Grade) in 2 gallons of water and apply at the rate of 2 gallons per 1000 square feet of surface. Apply first treatment as soon as bin is filled and leveled, but not later than the first week in October. Apply the second surface treatment 1 month later, followed by treatments at 2 month intervals. Use a piston type power sprayer equipped with agitator and nozzle capable of delivering a coarse spray. Repeat surface application 60 days later if warm temperatures prevail. Otherwise, delay further treatment until late winter.

Note — Both E. C. and WP are to be used in the full program.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.