

E-Z FLO SUCKER DOPE

AIDS IN THE CONTROL OF SUCKERS ON TOBACCO

CAUTION:
KEEP OUT OF THE REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

ACTIVE INGREDIENTS:

Aromatic petroleum derivative solvent

98.5%

INERT INGREDIENTS:

1.5%

100.0%

The manufacturer makes no warranties, express or implied, on using this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

SOLD BY

E-Z FLO CHEMICAL COMPANY

DIVISION OF GROWER SERVICE CORPORATION

ABERDEEN, NORTH CAROLINA

COLUMBUS, OHIO

LANSING, MICHIGAN

477-747

53-263-854

U.S.D.A. Reg. No. 635-332

DIRECTIONS FOR USE

Mix one gallon of this material with 3 to 4 gallons of water. Apply approximately 1 tablespoonful per plant at top of stalk, after topping.

LIQUID MEASURE EQUIVALENTS

5 teaspoons	1 tablespoon	16 tablespoons	1 cup
4 tablespoons	$\frac{1}{4}$ cup	2 tablespoons	1 liquid oz.
$5\frac{1}{3}$ tablespoons	$\frac{1}{3}$ cup	1 cup	$\frac{1}{2}$ pt
8 tablespoons	$\frac{1}{2}$ cup	2 cups	1 pt
$1\frac{2}{3}$ tablespoons	$\frac{2}{3}$ cup	4 cups	1 qt.
12 tablespoons	$\frac{3}{4}$ cup	4 qts	1 gal

1 part of any material per acre in 25 gallons of water = 1 table spoonfuls in 1 gallon water.

1 quart of any material per acre in 25 gallons of water = 25 table spoonfuls in 1 gallon water.

1 pint of any material per acre in 25 gallons of water = 12 table spoonfuls in 1 gallon water.

1 cup of any material per acre in 25 gallons of water = 25 table spoonfuls in 1 gallon water.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-FILL.

CAUTION

Harmful if swallowed.

Do not spray this material on the leaf surface since burning may result. If rain occurs soon after application the effectiveness may be reduced. This material is not for insect control!

E-Z FLO CLEAN SWEEP 50-W

CAUTION:
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS:

Diphenamid N, N-Dimethyl
 2,2-diphenylacetamide 50%

INERT INGREDIENTS: 50%

100%

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

[Faint, mostly illegible text providing detailed usage instructions for various crops and conditions.]

POTATOES (HARVEST): Apply 8-12 lbs. at planting or immediately after planting. Do not apply within 30 days of harvest.

POTATOES (SWEET): Apply 8-12 lbs. shortly after transplanting. Do not plant treated areas to crops not on this label.

STRAWBERRIES: Apply on established beds during fall season while strawberries are dormant or after summer harvest. A second application may be made 6 months later. Do not apply within 60 days of harvest. Do not use on variety Starbuck as injury may result.

TOBACCO: Apply 2-12 lbs. immediately after seeding or in small areas use 3-4.5 gal. per 1,000 sq. ft. or immediately after transplanting as overtop or directional spray. Two shallow (2 in.) cultivations (at second and lay-by) are suggested. Do not mix with other chemicals. Do not graze treated areas. Do not plant treated areas to crops not on this label.

WARNING

Harmful if swallowed. Induce vomiting and call a physician immediately. If spilled on hands or other skin areas, wash immediately with soap and water. In case of eye contact, flush with water for fifteen minutes and call a physician. Do not cover crop for food or feed within 6 months of last treatment.

