

TOBACCO FORMULA 2% PARATHION 4% THIODAN®



POISON



DANGER: KEEP OUT OF THE REACH OF CHILDREN

See Side Panel For Additional Warnings and Antidote Statements

ACTIVE INGREDIENTS:

Parathion (O, O-Diethyl O-p-nitrophenyl thiophosphate) 2.0%

Endosulfan (Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide) 4.0%

INERT INGREDIENTS: 94.0%

TOTAL 100.0%

*Thiodan is a registered trademark of Canadian Hoechst Limited.
U. S. Patent No. 2,799,685.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

Not Contents and Manufacturer
 On Container

4-14-71
 635-613

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

This "Tobacco Formula" Dust is formulated on a non-alkaline, sterilized tobacco base carrier. It is compounded especially for use on tobacco for the control of Aphids, Budworms, Cabbage Loopers, Hornworms, Cutworms, Vegetable Weevils, and certain other insects.
WARNING: Do not apply this product to tobacco beds with shaker cans or cloth bags. Applications in the field should be made with power dusting equipment only.

WARNING: No application should be made within 5 days of priming tobacco or 15 days of cutting. Avoid plant juices coming in contact with the skin of those persons engaged in cutting or priming the crop.

DIRECTIONS FOR USE

Insect	Dosage Per Acre Or As Otherwise Indicated	When And Where To Apply	Safety Restrictions
IN PLANT BED			
Aphids and Vegetable Weevils	0.4-0.5 lb/100 sq. yds.	On foliage at seven-day intervals beginning three weeks prior to transplanting for aphids or as needed for vegetable weevils.	Do not apply parathion within 5 days before transplanting. Should be applied only by a trained operator.
Cutworms	0.4-0.5 lb/100 sq. yds.	On foliage as needed.	
Garden Slugs and Pillbugs	0.4-0.5 lb/100 sq. yds.	These pests hide under boards, wall flaps and under debris in the plant bed. Applications should be directed at hiding places as needed.	
Insect	Dosage Per Acre Or As Otherwise Indicated	When And Where To Apply	Safety Restrictions
IN FIELD OR SHADE			
Aphids, Budworms, Hornworms and Cabbage Loopers	20-25 lbs.	On foliage at seven-day intervals beginning immediately following transplanting and continuing to end of priming season. Do not dust so as to obtain maximum coverage of the plant and particularly the bud for aphid and budworm control.	Should be applied only by a trained operator.

DANGER

KEEP AWAY FROM CHILDREN - This is a highly toxic insecticide. It is extremely poisonous to humans and animals. May be fatal if swallowed. Avoid contact with eyes, nose, mouth, and skin. Wash thoroughly with soap and water immediately after use. Do not eat, drink, or smoke while using this product. Wear a mask and protective clothing. Keep children and pets away from treated areas. Do not use in areas where food is stored. Do not use in areas where people or animals are present. Do not use in areas where children play. Do not use in areas where people or animals are present. Do not use in areas where children play.

FIRST AID TREATMENT - If swallowed, induce vomiting immediately. If inhaled, move to fresh air. If on skin, wash with soap and water. If in eyes, flush with water for 15 minutes.

INTERNAL - If swallowed, induce vomiting immediately. If inhaled, move to fresh air. If on skin, wash with soap and water. If in eyes, flush with water for 15 minutes.

SKIN - Wash immediately with soap and water. If irritation persists, seek medical attention.

EYES - Flush immediately with water for 15 minutes. Seek medical attention if irritation persists.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON

NOTE TO PHYSICIAN - Therapeutic doses of this insecticide are effective against certain insects. It is not intended for human use. It is extremely poisonous to humans and animals.

NOT FOR USE FOR STORAGE IN OR AROUND THE HOME

Always use this product in well-ventilated areas. Do not use in areas where people or animals are present. Do not use in areas where children play. Do not use in areas where people or animals are present. Do not use in areas where children play.


Decontamination and Disposal Instructions - Wash thoroughly with soap and water. Do not use in areas where people or animals are present. Do not use in areas where children play. Do not use in areas where people or animals are present. Do not use in areas where children play.

NET WEIGHT: 50 POUNDS

TOBACCO FORMULA 2% PARATHION 4% THIODAN



POISON



KEEP OUT OF REACH OF CHILDREN
 See Back Panel For Additional Warnings

ACTIVE INGREDIENTS:

Parathion (O, O-Diethyl O-p-nitrophenyl thiophosphate)	2.0%
Endosulfan (Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide)	4.0%
INERT INGREDIENTS:	94.0%
TOTAL	100.0%

*Thiodan is a registered trademark of Canadian Hoechst Limited
 U. S. Patent No. 2,799,685

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

4-14-71
635-613

On 10/10/71

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All treatments made concerning this product apply only when used as directed.

WARNING: Do not apply this product to tobacco plants with a maximum of 10% maturity. Application should be made with power-spraying equipment only.

WARNING: No application should be made within 10 days of harvest. Do not apply to plants that are being harvested or to plants that have been harvested.

DIRECTIONS FOR USE

Plant	Dosage Per Acre Or As Otherwise Indicated	When And Where To Apply	Safety Restrictions
A. PLANTED			
1. Tobacco	2.0% 4.0% 4.0% 4.0%	Apply at intervals of 7 to 10 days, beginning when plants are 10% mature and continuing until plants are 80% mature. Do not apply to plants that are being harvested or to plants that have been harvested.	Do not apply to plants that are being harvested or to plants that have been harvested.
2. Tobacco	2.0% 4.0% 4.0%	Apply at intervals of 7 to 10 days, beginning when plants are 10% mature and continuing until plants are 80% mature. Do not apply to plants that are being harvested or to plants that have been harvested.	Do not apply to plants that are being harvested or to plants that have been harvested.
B. FIELD OR SHADE			
1. Tobacco	2.0% 4.0% 4.0%	Apply at intervals of 7 to 10 days, beginning when plants are 10% mature and continuing until plants are 80% mature. Do not apply to plants that are being harvested or to plants that have been harvested.	Do not apply to plants that are being harvested or to plants that have been harvested.

NET WEIGHT: 50 POUNDS