

E-Z FLO PHOSDRIN* 4

(Contains 4 Lbs. Phosdrin Per Gallon)

ACCEPTED

DEC 14 1970
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 635-441 SUBJECT
TO ATTACHED COMMENTS.

DANGER: Keep out of the reach of children
SEE SIDE PANEL FOR ADDITIONAL WARNINGS AND ANTIPOISE STATEMENT



POISON



ACTIVE INGREDIENTS:	By Wt.
2-Carbomethoxy-1-Methylvinyl Dimethyl Phosphate, Alpha Isomer.....	28.1%
Related Compounds	18.7%
Xylene	45.8%
INERT INGREDIENTS:	7.4%
<hr/>	
*Shell Registered Trademark.	100.0%

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

DIRECTIONS FOR USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

CONTAINER DISPOSAL: BEFORE DISCARDING THIS CONTAINER, FILL IT WITH WATER, ADDING 3 TABLESPOONFULS CAUSTIC (ALKALINE) SUBSTANCE, SUCH AS WASHING SODA (SODIUM CARBONATE) OR LIME. ALLOW THE CONTAINER WITH WATER AND CAUSTIC TO STAND OVERNIGHT. BURY THE WASHINGS. EMPTY DRUMS MUST NOT BE USED IN ANY CONNECTION WITH FEED, FOOD OR DRINKING WATER. FOR 5 GALLON SIZE, WASH, PERFORATE AND BURY CONTAINER WHEN EMPTY. NEVER RE-USE.

Do not apply later than 1 day before harvest unless otherwise specified. Apply to give uniform coverage. Begin applications when insects first appear. Repeat as necessary. This product can be applied with conventional hydraulic sprayers or low volume ground applicators at 10 to 150 gals. per acre or with aerial equipment at 3 to 10 gals. per acre. Do not apply with hand sprayers.

ALFALFA, CLOVER: Aphids, Alfalfa Caterpillar — ¼ to ½ pint in water to cover 1 acre. Grasshoppers, Leafhoppers, Cutworms (Climbing), Red Spider Mites, Lygus Bugs, Alfalfa Weevil Larvae — ½ to 1 pint in water to cover 1 acre.

APPLES, PEACHES, PEARS, PLUMS, CHERRIES: Aphids, Red Spider Mites — ¼ to ½ pint per 100 gals. water. Grasshoppers, Lygus Bugs, Red-Banded Leaf Roller — ½ to ¾ pint per 100 gals. water. Do not apply to Cherries within 2 days of harvest. Do not apply more than 600 gals. per acre per application on cherries, peaches, and plums, nor more than 800 gals. per acre per application on apples and pears.

BEANS, PEAS, POTATOES, CUCUMBERS, TOMATOES: Aphids — ¼ to ½ pint in water to cover 1 acre. Grasshoppers, Leafhoppers, Red Spider Mites, Mexican Bean Beetle — (except potatoes) — ½ to 1 pint in water to cover 1 acre. **CAUTION:** Do not exceed ½ pint per acre on potatoes.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, COLLARDS, KALE, LETTUCE, SPINACH, TURNIPS, BEETS, CARROTS, CELERY, EGGPLANT, PEPPERS: Aphids — ¼ to ½ pint in water to cover 1 acre. Cabbage Looper, Imported Cabbage Worm, Grasshoppers, Leafhoppers, Salt Marsh Caterpillar, Red Spider Mites, Cutworm (Climbing), Dipterous Leafminer (Adults), Lygus Bugs, Corn Earworm, False Chinch Bug — ½ to 1 pint in water to cover 1 acre. Do not apply to Lettuce within 2 days of harvest; to Brussels Sprouts, Cauliflower, Celery, Collards, Beets, Kale and Turnips within 3 days and to Spinach within 4 days of harvest.

CANTALOUPE, HONEYDEW MELON, MUSKMELONS, WATERMELONS, SUMMER SQUASH: Aphids — ¼ to ½ pint in water to cover 1 acre. Cabbage Looper, Dipterous Leafminer (Adult), Leafhoppers, Lygus Bugs, Red Spider Mites, False Chinch Bug, Salt Marsh Caterpillars, Cutworms (Climbing), Grasshoppers — ½ to 1 pint in water to cover 1 acre.

GRAPES: Aphids — ½ to 1 pint in water to cover 1 acre. Leaf Folder, Leafhopper, Red Spider Mites, Red-Banded Leaf Roller, Lygus Bugs — 1 to 2 pints per acre. Do not apply within 2 days of harvest

ONIONS (Including Green Onions): Thrips, Cutworms (Climbing) — ½ to 1 pint in water to cover 1 acre.

RASPBERRIES: Aphids — ¼ to ½ pint per 100 gals. water. Red Spider Mites, Leafhoppers, Fruit Tree Leafroller — ½ to ¾ pint per 100 gals. water. Do not treat within 3 days of harvest.

NET CONTENTS OF CONTAINER

Do not exceed 1/2 pint per acre on potatoes.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, COLLARDS, KALE, LETTUCE, SPINACH, TURNIPS, BEETS, CARROTS, CELERY, EGGPLANT, PEPPERS: Aphids 1/4 to 1/2 pint in water to cover 1 acre. Cabbage Looper, Imported Cabbage Worm, Grasshoppers, Leafhoppers, Salt Marsh Caterpillar, Red Spider Mites, Cutworm (Climbing), Dipterous Leafminer (Adults), Lygus Bugs, Corn Earworm, False Chinch Bug 1/2 to 1 pint in water to cover 1 acre. Do not apply to lettuce within 2 days of harvest, to Brussels Sprouts, Cauliflower, Celery, Collards, Lettuce, Kale and Turnips within 3 days and to Spinach within 4 days of harvest.

CANTALOUPE, HONEYDEW MELON, MUSKMELONS, WATERMELONS, SUMMER SQUASH: Aphids 1/4 to 1/2 pint in water to cover 1 acre. Cabbage Looper, Dipterous Leafminer (Adult), Leafhoppers, Lygus Bug, Red Spider Mites, Imported Cabbage Worm, Salt Marsh Caterpillars, Cutworms (Climbing), Grasshoppers 1/2 to 1 pint in water to cover 1 acre.

GRAPES: Aphids 1/4 to 1/2 pint in water to cover 1 acre. Leaf Miner, Leafhopper, False Chinch Bug, Red Spider Mites, Leaf Roller, Lygus Bug 1/2 to 1 pint in water to cover 1 acre. Do not apply within 1 week of harvest.

ONIONS (Including Green Onions): True Vine Weevil, Onion Maggot 1/2 to 1 pint in water to cover 1 acre.

RASPBERRIES: Aphids 1/4 to 1/2 pint in water to cover 1 acre. Red Spider Mites, Imported Cabbage Worm, Salt Marsh Caterpillars, Cutworms (Climbing), Grasshoppers 1/2 to 1 pint in water to cover 1 acre.

Ones

13-293 810

13-293 810

DIRECTIONS FOR USE (Continued)

STRAWBERRIES: Aphids, Red Spider Mites, Trips 1/4 to 1/2 pint in water to cover 1 acre. Grasshoppers, Strawberry Leaf Roller, Salt Marsh Caterpillar, Lygus bugs 1/2 to 1 pint in water to cover 1 acre.

USE PRECAUTIONS

Do not apply under conditions favorable to the drift to food, forage or other plants if such drift is likely to cause them to become unfit for sale, use or consumption.

Toxic to fish and wildlife. Do not contaminate any body of water that apply to any area where such drift is likely to occur. Drift on treated areas may be killed.

This product is highly toxic to bees exposed to direct treatment. Do not apply to blooming crops or weeds which bees visit. For more information on bee protection, contact your Cooperative Agriculture Extension Office.

DESTROY CONTAINER WHEN EMPTY NEVER REUSE