

# E-Z FLO PARA-CAP

**DANGER**  
**KEEP OUT OF REACH OF CHILDREN**



**POISON**



See Side Panel for Additional Warnings and Antidote Statements

**ACTIVE INGREDIENTS:**

**Captan\*** ..... **25.0%**

**Parathion\*\*** ..... **7.0%**

**INERT INGREDIENTS:** ..... **68.0%**

**100.0%**

\*N-(trichloromethylthio)-4-cyclohexene-1,2-dicarboximide.

\*\*O,O-diethyl O-p-nitrophenyl thiophosphate.

**DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.**

**DIRECTIONS FOR USE**

Spray thoroughly for complete coverage. This product is not to be used around homes or home gardens.

**APPLES:** Scab, Frogeye Leaf Spot, Scale, and Aphids, use 4 lbs. per 100 gallons of water as required to maintain adequate coverage in the pre-bloom and blossom applications.

**Apple Shot-Hole Borer** — spray 4 lbs. per 100 gallons water when adults are active in May or September or after crop is harvested.

Do not use Parathion on McIntosh or related varieties. Do not apply within 14 days of harvest.

**PEACHES:** (NOT for use in California.) Plum Curculio, Oriental Fruit Moth, Plant Bugs, Aphid, San Jose, Forbes, Lecanium and Cottony Cushion Scale, Red Spider Mites, Peach Tree Borer, Lesser Peach Tree Borer, Brown Rot and Scab — use 4 lbs. per 100 gallons water. Repeat weekly beginning at petal fall for 2 or 3 applications, then at 10-day intervals for the cover sprays. Use full coverage spray for control of scale insects. Peach Tree Borer trunk spray — spray trunks of trees thoroughly during periods of Moth flight in July, August and September. Same dosage as above.

Do not apply to Peaches within 14 days of harvest. Do not apply more than 5 lbs. actual parathion per acre per year.

Do not use in combination with Lime, Sodium Hydroxide (Caustic), Lime Sulphur, Bordeaux Mixture, high lime Calcium Aresenate or any strongly alkaline material. Do not use in combination with or closely following oil sprays.

ACCEPTED

Apr 8, 1971

UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 FOR ECONOMIC POISON REGISTER-  
 ED UNDER NO. 635-623

**DANGER**

Poisonous if Swallowed, Inhaled, or Absorbed Through Skin! Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for Parathion protection. Rapidly Absorbed Through Skin! Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Do not use on D'Anjou Pears. Avoid use during flowering and pollination periods. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

**ANTIDOTE:** If Swallowed, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. **CALL A PHYSICIAN IMMEDIATELY! IF ON SKIN.** In case of contact, remove contaminated clothing and immediately wash skin with soap and water.

**PHYSICIAN'S NOTE:** Give atropine, preferably by injection, grains 1/100 two or three tablets at once and parenterally or orally every hour as required up to 30 tablets or until pupils dilate. 2-PAM is also antidotal and may be administered in conjunction with atropine. Never give morphine. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood tests.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crop. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOT FOR USE FOR STORAGE OF AEROSOLIZED PESTICIDES

STORE IN PLACE. DO NOT PLACE PRODUCT FROM EXHAUSTIVE TREATMENT BURN BAG IMMEDIATELY WHILE HOT. STAY OUT OF RAIN.

Net Contents and Manufacturer  
 On Container

# E-Z FLO PARA-CAP

KEEP OUT OF REACH OF CHILDREN



POISON



See Side Panel for Additional Warnings and Antidote Statements

ACCEPTED

UNDER THE FEDERAL INSECTICIDE ACT OF 1947 AND THE FEDERAL FOOD, DRUG AND COSMETIC ACT OF 1938 FOR ECONOMIC PESTICIDES REG. UNDER NO. 101-101-1

## ACTIVE INGREDIENTS:

**Captan\*** ..... 25.0%

**Parathion\*\*** ..... 7.0%

**INERT INGREDIENTS:** ..... 68.0%

**100.0%**

\*N-(trichloromethylthio)-4-cyclohexene-1,2-dicarboximide.  
 \*\*O,O-diethyl O-p-nitrophenyl thiophosphate.

**DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.**

## DIRECTIONS FOR USE

Spray thoroughly for complete coverage. This product is not to be used around homes or home gardens

**APPLES:** Scab, Frogeye Leaf Spot, Scale, and Aphids, use 4 lbs. per 100 gallons of water as required to maintain adequate coverage in the pre-bloom and blossom applications.

**Apple Shot-Hole Borer** — spray 4 lbs. per 100 gallons water when adults are active in May or September or after crop is harvested.

Do not use Parathion on McIntosh or related varieties. Do not apply within 14 days of harvest.

**PEACHES:** (NOT for use in California) Plum Curculio, Oriental Fruit Moth, Plant Bugs, Aphids, San Jose, Forbes, Lecanium and Cottony Cushion Scale, Red Spider Mites, Peach Tree Borer, Lesser Peach Tree Borer, Brown Rot and Scab— use 4 lbs. per 100 gallons water. Repeat weekly beginning at petal fall for 2 or 3 applications, then at 10-day intervals for the cover sprays. Use full coverage spray for control of scale insects. Peach Tree Borer trunk spray — spray trunks of trees thoroughly during periods of Moth flight in July, August and September. Same dosage as above.

Do not apply to Peaches within 14 days of harvest. Do not apply more than 5 lbs. actual parathion per acre per year.

Do not use in combination with Lime, Sodium Hydroxide (Caustic), Lime Sulphur, Bordeaux Mixture, high lime Calcium Aresenate or any strongly alkaline material. Do not use in combination with or closely following oil sprays.

### ANTIDOTE

### PHYSICIANS NOTE

Net Contents and Manufacturer On Container