

E-Z FLO M-S GARDEN DUST

WARNING:

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS:

| | |
|---------------------------------------|------|
| *Malathion | 3.0% |
| Carbaryl (1-Naphthyl methylcarbamate) | 2.5% |

| | |
|--------------------|--------|
| INERT INGREDIENTS: | 94.5% |
| | <hr/> |
| | 100.0% |

*O, O-Dimethyl Dithiophosphate of Diethyl mercaptosuccinate.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

Begin applications when insects or their damage is first noticed. Repeat applications on a 5 to 7 day schedule or as necessary for control. Applications should be made with a good dust gun or blower to give coverage to both upper and lower surface of the leaves. Apply on a per acre basis the rates as indicated below.

BEANS: For the control of Japanese Beetle, Velvet Bean Caterpillar, Flea Beetle, Western Bean Cutworm, Bean Leaf Beetle, Mexican Bean Beetle, Leafhoppers, Aphids and Cucumber Beetle. Apply 40 to 47 lbs. For the control of Corn Earworm, Armyworms, Stink Bugs, and Tarnished Plant Bug — Apply 40 to 58 lbs. Do not apply within 1 day of harvest.

CABBAGE AND COLLARDS: For the control of Harlequin Bug, Flea Beetles, Leafhoppers, and Aphids — Apply 40 lbs. For the control of Six Spotted Leafhopper — Apply 40 to 60 lbs. For the control of Corn Earworm, Armyworm, Tarnished Plant Bug, and Stink Bug — Use 40 to 80 lbs. Do not apply to cabbage within 7 days nor to collards within 14 days of harvest.

CUCUMBERS: For the control of Cucumber Beetle, Squash Bug, Aphids, Flea Beetle, Spider Mites, Melonworm, Pickleworm, and Leafhoppers — Apply 40 lbs. Do not apply unless the plants are dry. Do not apply within 1 day of harvest.

EGGPLANT: For the control of Colorado Potato Beetle, Flea Beetle, and Leafhoppers — Apply 20 to 40 lbs. For the control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Stink Bugs, Lace Bugs, and Tarnished Plant Bugs — Apply 40 to 80 lbs. Do not apply within 3 days of harvest.

PEAS: For the control of Leafhoppers, and Colorado Potato Beetle — Apply 40 lbs. For the control of Armyworms and Grasshoppers — Apply 40 to 60 lbs. Do not apply within 3 days of harvest. Do not forage or feed vines to livestock.

PEPPERS: To control Leafhoppers, Colorado Potato Beetle, and Flea Beetle — Apply 20 to 40 lbs. For the control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Stink Bug, Lace Bug, and Tarnished Plant Bug — Apply 40 to 50 lbs. For Pepper Maggot control — Apply 50 lbs. Do not apply within 3 days of harvest.

POTATOES (Irish): For the control of Colorado Potato Beetle, Flea Beetle, and Leafhoppers — Apply 20 to 40 lbs. For the control of Aphids, Tomato Fruit Worm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Stink Bug, Lace Bug and Tarnished Plant Bug — Apply 40 to 50 lbs. For the control of False Chinch Bug and Mealy Bugs — Apply 50 lbs. Apply as a foliage application only.

MELONS: For the control of Pickleworm and Melonworm — Apply 20 to 40 lbs. For the control of Cucumber Beetles, Squash Bug, Flea Beetles, Leafhoppers, Aphids, and Spider Mites — Apply 40 lbs. Do not apply within 1 day of harvest.

OKRA: For the control of Aphids — Apply 27 to 40 lbs. For the control of Japanese Beetle — Apply 40 lbs. For the control of Corn Earworm and Stink Bug — Apply 40 to 50 lbs. Do not apply within 1 day of harvest.

PUMPKINS: For the control of Pickleworm and Melonworm — Apply 20 to 40 lbs. For the control of Aphids, Spider Mites, Leafhoppers, Cucumber Beetles, Squash Bugs, and Flea Beetles — Apply 40 lbs. Do not apply within 3 days of harvest.

SQUASH: For the control of Pickleworm and Melonworm — Apply 20 to 40 lbs. For the control of Aphids, Spider Mites, Cucumber Beetle, Squash Bug, Flea Beetle, and Leafhoppers — Apply 40 lbs. Do not apply within 1 day of harvest.

TOMATOES: For the control of Colorado Potato Beetle, Flea Beetle, and Leafhoppers — Apply 20 to 40 lbs. For the control of Aphids, Spider Mites, and Tomato Russet Mites — Apply 47 to 60 lbs. For the control of Drosophila — Apply 53 lbs. For the control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Stink Bug, Lace Bug and Tarnished Plant Bug — Apply 40 to 80 lbs. Do not apply within 1 day of harvest, unless application exceeded 66 lbs., then do not apply within 5 days of harvest.

CAUTION

Harmful if inhaled or swallowed. Avoid breathing of dust. Do not use internally. Skin contact may be harmful. Avoid contact. Wash hands and face after use. Do not eat, drink, or smoke, or use tobacco or bath after using.

Do not apply when weather conditions favor drift from treated areas.

Avoid contamination of feed and foodstuffs.

Highly toxic to fish. Keep out of lakes, ponds, and streams.

This product is highly toxic to bees exposed to direct treatment or residues. For more complete information may be obtained from your cooperative Agricultural Extension Service.

NOTE TO PHYSICIAN: Carbaryl is a moderate to variable central nervous system inhibitor. Atropine or other anticholinergic agents should be administered prophylactically. Careless use may cause the moderate to severe atropine-like symptoms.

Not Contain... Manufacturer

