

E-Z FLO CYPREX[®], PARATHION, SULPHUR 2-2-20 DUST

DANGER:
KEEP OUT OF THE REACH OF CHILDREN



POISON



635-626
ACCEPTED
Dec 13 1970
UNDER THE FEDERAL INSECTICIDE
AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO.

USE PRECAUTIONS

Do not prune fruit trees or thin fruit within 15 days after application. Do not apply during bloom when honey bees or other pollinators are active. Avoid contact with dust drift on flowers, shrubs, or fruits, vegetables, and forage crops. Keep children and animals off treated areas until chemical has been washed into the soil by sprinkling or rain and the grass is dry.

Do not re-use container. Destroy by burying in a safe place. Do not burn.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates and do not apply closer to harvest than specified. Consult your State Agricultural Experiment Station for specific information consistent with labeling as the timing, number and rate of applications may vary with local conditions.

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

ACTIVE INGREDIENTS:

Dodine (n-Dodecylguanidine acetate) 2.0%
Parathion (O,O-Diethyl O-p-Nitrophenyl Thiophosphate) 2.0%
Sulphur 20.0%

INERT INGREDIENTS:

76.0%
100.0%

*Cyprex is a registered trademark of American Cyanamid Company. U. S. Patent No. 2,867,562.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

APPLES: For the control of Scab, Powdery Mildew, Codling Moth, Plum Curculio, Orange Tortrix, San Jose Scale, Forbes Scale, Scurfy Scale, Red Banded Leaf Roller, Apple Maggot, Woolly Apple Aphid, Bud Moth, Fruit Tree Leaf Roller, Rosy Apple Aphid, Green Apple Aphid, Leafhopper, European Red Mite and Two-spotted Mite. Apply 50-65 lbs. per acre at 7 day intervals or as needed to maintain control from pre-bloom period through 1st cover. Repeat at 7-10 day intervals throughout the summer months. Make full coverage applications. Do not apply within 14 days of harvest. Do not apply to susceptible varieties such as McIntosh, Cortland, and related varieties, Golden Delicious, Grimes Golden, Rhode Island Greening or Jonathan. Do not use treated apples in the manufacture of apple pomace for uses in livestock feeds.

CHERRIES: For the control of Leaf Spot, Brown Rot, Aphids, Mites, Leaf-miner, Plum Curculio, Cherry Fruit Worm, Oriental Fruit Moth, Bud Moth, Tortrix, Cankerworm, San Jose Scale. Apply 50-65 lbs. per acre beginning at petal fall or 5 days after first fruit firmness. Repeat every 7-10 days and immediately after every rain. Make full coverage application. Do not apply within 14 days of harvest. Make a post harvest application for additional protection.

Net Contents and Manufacturer
On Container

DANGER

Extremely hazardous if swallowed, inhaled or absorbed through the skin. Avoid getting it on the skin, in the eyes, or on clothing. Rapidly absorbed through skin. Do not breathe dust. Wear a respirator which has been tested by the U. S. Department of Agriculture for parathion protection. Wear waterproof or protective clothing, natural rubber gloves, hat, and goggles. Keep away from all feed and food products. Wash hands, arms and face with soap and water thoroughly after application or before eating or smoking. Wash contaminated clothing with hot water and soap before reusing. Keep all unprotected persons or animals out of the operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticides and volatile residue has dissipated.

FIRST AID TREATMENT: Call a doctor (note for doctor): Therapeutic doses of atropine are effective as an antidote. 2-PAM is also antidotal and may be administered in conjunction with atropine. **INTERNAL:** If swallowed, make patient vomit by giving warm salt water or soapy water. Administer milk or water freely. Have patient lie down and keep warm. Never give anything by mouth to an unconscious patient. **SKIN:** Immediately remove all contaminated clothing and flush skin with water for at least fifteen minutes. **EYES:** Immediately flush with water for at least fifteen minutes. **INHALATION:** Give two atropine tablets at once if patient shows symptoms of blurred vision or pin point pupils.

E-Z FLO CYPREX[®], PARATHION, SULPHUR 2-2-20 DUST

DANGER:

KEEP OUT OF THE REACH OF CHILDREN



POISON



635-626
Rec. 13 1970
REGISTER.

ACTIVE INGREDIENTS:

Dodine (n-Dodecylguanidine acetate)	2.0%
Parathion (O,O-Diethyl O-p-Nitrophenyl Thiophosphate)	2.0%
Sulphur	20.0%

INERT INGREDIENTS:	76.0%
	100.0%

*Cyprex is a registered trademark of American Cyanamid Company. U. S. Patent No. 2,867,562.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

APPLES: For the control of Scab, Powdery Mildew, Codling Moth, Plum Curculio, Orange Tortrix, San Jose Scale, Forbes Scale, Scurfy Scale, Red Banded Leaf Roller, Apple Maggot, Woolly Apple Aphid, Bud Moth, Fruit Tree Leaf Roller, Rosy Apple Aphid, Green Apple Aphid, Leafhopper, European Red Mite and Two-spotted Mite. Apply 50-65 lbs. per acre at 7 day intervals or as needed to maintain control from pre-bloom period through 1st cover. Repeat at 7-10 day intervals throughout the summer months. Make full coverage applications. Do not apply within 14 days of harvest. Do not apply to susceptible varieties such as McIntosh, Cortland, and related varieties, Golden Delicious, Grimes Golden, Rhode Island Greening or Jonathan. Do not use treated apples in the manufacture of apple pomace for uses in livestock feeds.

CHERRIES: For the control of Leaf Spot, Brown Rot, Aphids, Mites, Leaf miner, Plum Curculio, Cherry Fruit Worm, Oriental Fruit Moth, Bud Moth, Tortrix, Cankerworm, San Jose Scale. Apply 50-65 lbs. per acre beginning at petal fall or 5 days after first fruit fly emerges. Repeat every 7-10 days and immediately after every rain. Make full coverage application. Do not apply within 14 days of harvest. Make a post harvest application for additional protection.

USE PRECAUTIONS

Do not prune fruit trees or thin fruit within 15 days after application. Do not apply during bloom when honey bees or other pollinators are active. Avoid contact with dust drift on flowers, shrubs, or fruits, vegetables, and forage crops. Keep children and animals off treated areas until chemical has been washed into the soil by sprinkling or rain and the grass is dry.

Do not re-use container. Destroy by burying in a safe place. Do not burn.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates and do not apply closer to harvest than specified. Consult your State Agricultural Experiment Station for specific information consistent with labeling as the timing, number and rate of applications may vary with local conditions.

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DANGER

Net Contents and Manufacturer On Container