

E-Z FLO POLYRAM® 5 BORON DUST (WITH BORAX)

FUNGICIDE-NUTRITIONAL

CAUTION:
KEEP OUT OF THE REACH OF CHILDREN

See side panel for additional cautions.

DIRECTIONS FOR USE

PEANUTS - For the control of Cercospora Leaf Spot and as an aid in correcting the deficiency of boron. Apply undiluted as a dust at the rate of 20 pounds per acre when spots first appear on old leaves at the base of the plants. Only one application should be made per season. Dusting should be done with a machine capable of depositing a thin uniform coating of dust on all above ground plant surfaces. Dusting should only be done when the foliage is dry. Severe burning may result if application is made when the leaves are wet. Use of more than the recommended dosage may result in reduced yields.

ACTIVE INGREDIENTS:

A mixture of 5.2 parts by weight (83.9%) of ammoniates of [ethylenebis(dithiocarbamate)] zinc* with 1 part by weight (16.1%) ethylenebis [dithiocarbamic acid] bimolecular and trimolecular cyclic anhydrosulfides and disulfides..... 5%


INERT INGREDIENTS: 95%

100%

*Boron expressed as metallic..... 2.52%
 Zinc expressed as metallic..... 0.87%

®Polyram is a registered Trademark of Badische Anilin & Soda Fabrik A. B.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.



Net Contents and Manufacturer On Container

CAUTION

Keep out of the reach of children. May be harmful if swallowed or inhaled. Do not breathe dust. May cause irritation of eyes, nose, throat and skin. In case of contact flush with plenty of water; for eyes, get medical attention. Avoid contamination of feed and foodstuffs.

Do not feed treated forage to dairy animals being finished for slaughter.

Dispose of wastes by burying in non-crop lands away from water supplies. Bags should be disposed of by burying with wastes or by burning. Keep out of smoke.