

E-Z FLO POTATO-TOMATO 7-5 DUST

CAUTION:
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS:

Copper (In Basic Copper Sulphate)	
Expressed As Metallic.....	7.00%
Carbaryl (1-naphthyl N-methylcarbamate).....	5.00%
INERT INGREDIENTS:.....	88.00%
Total	100.00%

® Trade Mark of Union Carbide Corporation for the active ingredient carbaryl insecticide. U.S. Pat. Nos. 2,903,478; 3,009,855

CAUTION


Harmful if inhaled or swallowed. Avoid breathing of dust. Do not take internally. Skin contact may be harmful.

Continued on side/back panel-

512-9

EPA Est. 635-MI-1; OH-1

E. P. A. Reg. No. 635-629



DIRECTIONS FOR USE

CUCUMBERS, MELONS, PUMPKINS, SQUASH, and WATERMELONS: For the control of downy mildew, anthracnose, scab, *Macrosporium* leaf spot, angular leaf spot, pickleworm, melonworm, cucumber beetle, flea beetle, leafhopper and squash bug. Apply 20 lbs. per acre. To avoid possible injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days. Sevin injures watermelons in Florida.

POTATOES: For the control of early blight, late blight, Colorado potato beetle, flea beetle, leafhopper, European corn borer, fall armyworm, lace bug, lygus bug, stink bug, tomato fruitworm, tomato hornworm, and tarnished plant bug. Apply 20 to 40 lbs. per acre.

TOMATOES and PEPPERS: For the control of early blight, late blight, *Septoria* leaf spot, Colorado potato beetle, flea beetle, leafhopper, European corn borer, fall armyworm, stink bug, tomato fruitworm, tomato hornworm, tarnished plant bug, and climbing cutworm. Apply 20-40 lbs. per acre. Use high rate for climbing cutworm control.

CAUTION

Continued from front panel-

Avoid contact. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

Avoid Contamination of Food, Feed, Feeding Troughs and Watering Receptacles.

This product is toxic to fish. Keep out of lakes, ponds, and streams. Do not apply to any area not specified on the label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Carbaryl can be injurious to tender foliage. Do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days. Local weather and other conditions may affect the use of this material. Consult the State Agricultural Experiment Station for additional information consistent with label directions and cautions.

Do not re-use container. Destroy when empty.