

E-Z FLO CHLORDANE 5 DUST

CAUTION:

KEEP OUT OF THE REACH OF CHILDREN

635-642

ACTIVE INGREDIENTS:

Technical Chlordane

(Equivalent to 3% Octachloro-4,7-methanotetrahydroindane and 2% related compounds)

INERT INGREDIENTS:

ACCEPTED

NOV 30 1972

51.0% FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 99507 SUBJECT
TO GENERAL COMMENTS

100.0%

Net Contents and Manufacture
On Container

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

USE ONLY AS DIRECTED: Apply at the rate of 25 to 40 pounds per acre unless otherwise indicated.

STRAWBERRIES — Spittle bug. Apply 2 weeks before blossoming. Do not apply after fruit begins to form.

ONIONS — Control maggots and thrips. For maggot control, apply dust around base of the plants as soon as they are set in the field. Wash into soil. Apply only at planting time. Do not apply to green or spring onions.

APPLES — Bagworms, Grasshoppers, codling moth, oriental fruit moth, and Plum curculio. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Do not apply within 30 days of harvest.

BEANS — Flea beetles and Green stink bugs. Do not apply to green or snap beans after pods begin to form.

LETTUCE — Armyworm, Climbing cutworm, Lygus bug, and Serpentine leafminer. Do not apply after edible parts start to form.

POTATOES — Blister beetle, Colorado potato beetle, Flea beetle, adults and larvae, Mole crickets, Serpentine leafminer, and White fringed beetle. Apply at the rate of 25 to 30 pounds per acre. As a preplant soil treatment, for control of white grubs, flea beetle larvae, and wireworms, apply at the rate of 50 pounds per acre. at the rate of 50 pounds per acre.

TOMATOES — Leafminer. Do not apply after fruit begins to form.

CAUTION

Harmful if swallowed or inhaled. Contact with skin can cause toxic symptoms. Avoid breathing dust mist. In case of contact with skin wash with soap and water. Avoid contamination of feed or foodstuffs.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, ponds, and streams. Do not apply to any area not specified on the label. Do not apply when weather conditions favor drift of dust from the crop areas treated. Birds feeding on treated area may be killed.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.