

# E-Z FLO Prime Leaf Tobacco Spray Insecticide



Net Contents 1 Gallon  
(Contains 1 Lb. of Thiodan and 1 Lb. Malathion Per Gallon)

**ACCEPTED**  
 SEP 23 1975  
 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, by the pesticide regulatory board.  
 EPA Reg. No. 635-646

## WARNING: KEEP OUT OF REACH OF CHILDREN

See side panels for additional warnings.

ACTIVE INGREDIENTS:	
*Endosulfan (Hexachlorocyclohexane-2,4,3-benzothiazole epoxide)	12.20%
**Malathion	12.20%
Xylene	70.95%
INERT INGREDIENTS	4.65%
	100.00%

\*Thiodan is a registered trademark of Canadian Hoechst Limited.  
 \*\*O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate.  
 The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

56-17

EPA Est. 635-MI-1, OH-1

E.P.A. Reg. No. 635-646

## SOLD BY E-Z FLO CHEMICAL COMPANY

Division of Kirsto Company  
 Lansing, Michigan • Columbus, Ohio

## WARNING

Hazardous if swallowed, inhaled or absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water. For eyes, get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking, wear clean clothing. During commercial or prolonged exposure in spray-mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type passed by the U.S. Bureau of Mines for endosulfan protection. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Do not contaminate or store near feed or foodstuffs.

**NOTE TO PHYSICIANS:** Malathion may cause cholinesterase inhibition. Atropine is antidotal; 2-PAM is also antidotal and may be administered in conjunction with atropine. Thiodan produces central nervous system excitation which responds to the barbituric acid derivatives.

Continued on panel to right-

## WARNING (Continued)

**FLAMMABLE:** Keep away from heat and open flame. Keep container closed. Leaking packages should be removed to safe place.

Do not store below 20 degrees F.

This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Do not apply where it is not likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Rinse equipment and containers and dispose of wastes by burying at least 18" deep in non-crop areas away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

Do not allow this material to drift onto neighboring crop or non-crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues, or other undesirable results may occur.

Destroy Container When Empty, Never Re-Use.

## DIRECTIONS FOR USE

**TOBACCO: PLANT BEDS-** Aphids, Cabbage Looper, Tobacco Budworm, Tobacco Hornworm, Tobacco Flea Beetle- Use 4 qts. per 100 gals. of water. Apply 3 gals. of finished spray per 100 sq. yards seed bed.  
**FIELDS-** Aphids, Cabbage Looper, Tobacco Budworm, Tobacco Flea Beetle, Tobacco Hornworm- Use 2 to 4 qts. Stink Bugs- Use 2 to 4 qts. Grasshoppers- Use 4 qts. Dosage rates given should be used in sufficient water to provide uniform coverage of one acre.

Application of Prime Leaf Tobacco Spray should begin when insects first appear and repeat at 3 to 5 day intervals depending upon the severity of the infestation and local conditions. Full coverage of plants during application is essential for good control. To mix, fill spray tank to 1/2 full of water with agitator running, add the required amount of this material, then fill the spray tank to the rest of the tank with water. When applying this material by aircraft, mix the recommended amount with sufficient water to provide a minimum of 2 gals. of finished spray per acre.