

# E-Z FLO NEW IMPROVED ROSE GUARD

## CAUTION:

**KEEP OUT OF THE REACH OF CHILDREN**

### ACTIVE INGREDIENTS:

Captan (N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide) .....	10.00%
Carbaryl (1-Naphthyl methylcarbamate)* .....	10.00%
Malathion (O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate) .....	6.25%
2-(1-Methylheptyl)-4,6-dinitrophenyl crotonate and other nitrophenols and derivatives, chiefly 1-methylheptyl-4,6-dinitrophenol** .....	1.25%
<b>INERT INGREDIENTS:</b> .....	<b>72.50%</b>
<b>Total</b>	<b>100.00%</b>

\*Trade Mark of Union Carbide Corporation.  
U.S. Pat. Nos. 2,903,478; 3,009,855.

\*\*Karathane.

**DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE**

## GENERAL DIRECTIONS

Make first application before diseases become established or when insects first appear. Repeat applications every 7 to 10 days. Do not apply if temperature is over 90° F.

## PREPARATION OF SPRAY

Mix with water at the rate of 10 lbs. per 100 gals. water per acre (1.6 oz. per 1 gal. water per 435 sq. ft.). Use a pressure sprayer with agitation. Spray thoroughly, applying to all plant surfaces.

## USE AS A DUST

Apply lightly and evenly to all plant parts at the rate of 10 lbs. per acre (3.7 oz. per 1,000 sq. ft.).

**ROSES:** For the control of black spot, powdery mildew, aphids, spider mites, rose slug, lace bugs, Japanese beetle, flea beetles, plant bugs, leafhoppers, thrips, mealy bugs and tarnished plant bugs.

**FLOWERS:** For the control of powdery mildew, aphids, plant bugs, spider mites, lace bugs, Japanese beetle, flea beetles, leafhoppers, thrips, mealy bugs and tarnished plant bugs.

## CAUTION

**CAUTION:** Avoid inhalation of dust or spray mist. Harmful if swallowed or inhaled. May cause skin irritation. Avoid contact with skin; wash thoroughly after using. Do not get in eyes. Avoid contamination of feed or foodstuffs. Keep out of reach of children.

Toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

ACCEPTED