

[635-653]

E-Z FLO PYRETHRIN SPRAY

CAUTION: Keep out of the reach of children.
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

ACTIVE INGREDIENTS:	
Piperonyl Butoxide Technical*	60.0%
Pyrethrins	6.0%
Petroleum Distillates	18.0%
INERT INGREDIENTS:	
	18.0%
TOTAL:	100.0%

*Equivalent to 48.0% (Butylcarbityl) (6-propylpiperonyl) Ether and 12.0% of other related compounds

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

DIRECTIONS FOR USE

E-Z Flo Pyrethrin Spray is designed for use alone as a clean-up spray or as a pre-harvest spray where other materials cannot be used because of residue restrictions. When applied according to accepted use cautions and directions, Pyrethrin Spray may also be used in combination with other insecticides where resistant insecticides are a problem, or as an "exciter" while insects are flushed out of hiding and into contact with spray residues.

Apply the listed amount per acre and repeat as required to maintain effective control. Use in sufficient water for thorough coverage (usually 100 gallons) unless otherwise noted, for listed crops. Coverage of upper and lower leaf surfaces is essential for good control.

When used alone, E-Z Flo Pyrethrin may be used up to and including day of harvest.

ASPARAGUS: Asparagus Beetle — 2 to 6 oz.

BEANS: Aphids, Leafhoppers, White Fly, Mexican Bean Beetle, 12-Spotted Cucumber Beetle — 2 to 6 oz.

BROCCOLI, CABBAGE, BRUSSELS SPROUTS, CAULIFLOWER: Aphids, Cabbage Looper, Cross-Stripped Cabbageworm, Diamondback Moth Larvae, Flea Beetle, Harlequin Bug, Imported Cabbageworm, Stink Bug — 2 to 6 oz.

CELERY: Green Peach Aphid, Cabbage Looper, Leaf Tier — 2 to 6 oz.

CRANBERRIES: Fireworms, Leafhopper — 2 to 6 oz.

EGGPLANTS: Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid — 2 to 6 oz.

LETTUCE: Cabbage Looper, Green Peach Aphid, Diamondback Moth Larvae, Imported Cabbageworm — 2 to 6 oz.

PEPPERS: Green Peach Aphid — 2 to 6 oz.

POTATOES: Aphids, Colorado Potato Beetle, Flea Beetle, Leafhopper — 2 to 6 oz.

RADISH: Aphids, Flea Beetle — 2 to 6 oz.

SPINACH: Aphids, Cabbage Looper, Webworm — 2 to 6 oz.

TOMATOES: Green Peach Aphid, Colorado Potato Beetle, Stink Bug, Flea Beetle — 2 to 6 oz.

MOSQUITOES: May be used to control mosquitoes in areas where they breed. To protect agricultural workers during harvest, use at 1 oz. per 4 gallons of water. Apply by fixed wing aircraft.

DROSOPHILA: Fruit Fly Control. Dilute at the rate of 1 part Pyrethrin Spray with 1200 parts water (1 pint per 150 gallons or 1 tablespoonful with 4 gallons water). Thoroughly mix the emulsion in the spray tank. Make treatment as follows:

12-22

53-263-814

U.S.D.A. Reg. No. 635-653

DIRECTIONS FOR USE (Continued)

1. Apply liberally to tomatoes and other fruit in the field. Use sprayers at a high pressure for applying at the rate of 1 gallon per 100 square feet of a two-ton load of tomatoes. Direct the spray for a maximum coverage of the fruit. It is important to spray between and beneath the containers. This spray is not a contact insecticide, but the emulsion loosens any dead flies so that they are readily washed from the fruit.

2. Spray the raw stock stacked in the yard.

3. Dip baskets in the diluted spray after dumping the fruit to kill adhering larvae and pupae.

4. After washing and cleaning up the inside of the processing plant and just prior to bringing fruit into it, the entire space inside the cannery should be sprayed at a dilution of 1 part Pyrethrin Spray to 59 parts water (1 pint with 7 gallons 3 pints water or 2 tablespoonfuls with 3 1/4 pints water). At the rate of 1 gallon to 750 square feet (1 quart to 187 sq. ft.) direct spray on walls, ceilings and floors paying special attention to forcing the spray into all cracks and crevices for the control of ants, roaches, silverfish, crickets, spiders and cheese mites.

For use as a space spray in the processing plants, this dilution will give excellent control of flying insects, i.e. fruit flies, house flies, hornets, grain moths, gnats, mosquitoes and skipper flies. Apply at the rate of one-half ounce per 1000 cubic feet of space. Do not spray while the plant is in operation, as dead flies may fall into containers of the product being processed. Do not remain in treated areas and ventilate the area after treatment is completed.

SPACE SPRAY: To control flying insects such as fruit flies, hornets, wasps, grain moths, gnats, mosquitoes and skipper flies, dilute one part Pyrethrin Spray with 59 parts water (2 tablespoonfuls with 3 1/4 pints water). Use at the rate of one-half ounce per 1000 cubic feet of space. Direct the space treatment upward and whenever practical keep doors and windows closed for at least ten minutes after application. Do not remain in treated areas and ventilate the area after treatment is completed.

CAUTION

HARMFUL IF SWALLOWED.

USAGE CAUTIONS: Do Not Use in Undiluted Form.

This product is toxic to fish. Keep out of lakes, streams or ponds. Apply this product only as specified on this label.

Do not re-use empty containers. Destroy by perforating or crushing and bury in a safe place.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE

ACCEPTED

APR 29 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROBOTICIDE ACT
FOR ECONOMIC POISON REGISTER
NO. UNDER NO. 635-653