

E-Z FLO 10% SEVIN* GRANULES

FOR THE CONTROL OF EUROPEAN CORN BORER

CAUTION:
KEEP OUT OF THE REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) 10%


INERT INGREDIENTS: 90%

100%

U. S. Pat. No. 2,903,478

*Trade mark of Union Carbide Corporation for the active ingredient 1-naphthyl N-methylcarbamate (carbaryl)

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE



DIRECTIONS FOR USE

CORN: For the Control of European Corn Borer: Use 15 to 20 pounds per acre for first-generation borer larvae in whorls. Use suitable ground or air equipment to insure good coverage. Apply when pests appear in damaging numbers and repeat 7 to 14 days later if necessary.

Tolerances established under the Federal Food, Drug, and Cosmetic Act permit the sale of corn and corn forage bearing probable SEVIN residues. When SEVIN is used in accordance with label directions, corn or corn forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in meat or milk.

CAUTION

Harmful if inhaled or swallowed. Avoid breathing of dust. Do not take internally. Avoid contact with skin. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Avoid contamination of food, feed, water supplies, streams and ponds.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.