

Net Contents and Manufacture
On Container

635-659
ACCEPTED
DEC 19 1972
UP
PUN
FO
ED II
TO ALL

E-Z-FLO KOCIDE*-THIODAN® DUST

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENTS:

Endosulfan (hexachlorohexahydromethano-
2, 4, 3-benzodioxathiepin oxide)

3.00%

Copper hydroxide*

6.74%

INERT INGREDIENTS:

90.26%

Total

100.00%

DIRECTIONS FOR USE

This dust is formulated to control certain insects and diseases listed below. In general, applications should be made at 7 to 14 day intervals, depending on weather conditions and the severity of insect or disease problems. For disease control, applications should be started before diseases occur. Local weather and other conditions may affect the use of this material. State agricultural experiment stations may issue recommendations to fit local conditions.

*Reg. TM Kennecott Copper Corporation

*Thiodan is a registered trade mark of Canadian Hoechst Limited.
U.S. Patent No. 2,799,685.

*Metallic Copper Equivalent—4.39%
Manufactured under U.S. Patent No. RE. 24324

POTATOES: For control of aphids, Colorado Potato Beetle, Leafhopper Flea Beetle, Southern Armyworm, Green Stink Bug, Leaf-footed Plant bugs, Potato Tuberworms, and Whitefly; for the Control of Early and Late Blight of Potatoes: Apply this dust at the rate of 33 pounds per acre at 7 to 14 day intervals, depending on weather conditions and insect and disease incidence. Under conditions which favor Blight development, more frequent applications are required than during hot weather. Begin dusting soon after potato plants show above ground; continue throughout the growing season. Thorough coverage is important. No time limitations. Do not plant root crops other than carrots, potatoes, and sweet potatoes as follow-up crops.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing of dust. Avoid contact with skin, eyes or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Avoid contamination of feed and foodstuffs. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

TOMATOES: For control of Aphids, Colorado Potato Beetle, Flea Beetle, Whitefly, Green Stink Bug, Blister Beetle and Early Blight. Apply at the rate of 33 pounds per acre. Do not apply within 1 day of harvest.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds, and streams. Do not apply in any manner not specified on the label. Do not apply when weather conditions favor drift of dust from crop areas treated.

CUCUMBERS: For the control of Downy Mildew, Angular leaf spot, Aphids, Cucumber Beetles, Striped Cabbage Flea Beetle and the Squash Vine Borer. Apply at the rate of 33 pounds per acre when plants are two to six inches high and continue as needed.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

SUGAR BEETS: For control of Cercospora Leaf Spot, Green Peach Aphid and the Sugar Beet Webworm. Apply at the rate of 25 to 33 pounds per acre. Apply at 10 day intervals as needed. Do not feed treated tops to livestock. Do not apply within 30 days of harvest.

Do not reuse container. Destroy when empty.

210-31

53-261-625

E.P.A. Reg. No. 635-659