

E-Z FLO 2,4-D SODIUM SALT

CAUTION:
KEEP OUT OF THE REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

ACTIVE INGREDIENT:

**2,4-Dichlorophenoxyacetic acid,
sodium salt, monohydrate***

INERT INGREDIENTS:

5.6%

100.0%

*Equivalent to 80 % 2,4-Dichlorophenoxyacetic acid.

The manufacturer makes no warranties, express or implied, concerning the product or its use which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

DIRECTIONS FOR USE

WEED CONTROL IN ASPARAGUS: To control wild morning glory and certain other broad-leaved weeds in asparagus plantings, use 2 to 2 1/2 pounds per acre in approximately 60 gallons of water. In airplane application, use 2 pounds in approximately 12 gallons of water per acre. A wetting agent can be added if desired in both ground and airplane sprays. Apply on actively growing weeds, usually in April or May. Make no more than 2 applications at least 1 month apart during the harvest season. Post-harvest spraying should be only by ground rig, using drop nozzles to avoid spraying the fern. For young seedlings, do not exceed 1.25 lbs per acre. Be sure that use of this product conforms to all applicable regulations.

Local conditions may affect the use of this chemical. Consult State agricultural extension or Experiment Station weed specialists for specific recommendations for local weed problems and for information on possible lower dosages.

CAUTION

CAUTION: Irritating Dust. Avoid inhaling the Dust. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Do not store near seeds, insecticides, fungicides or fertilizers.

Do not apply 2,4-D Sodium Salt directly to, or otherwise permit it to come in contact with, vegetables, flowers, grapes, fruit trees, ornamentals, cotton, or other desirable plants which are sensitive to 2,4-D Sodium Salt. Do not permit spray mist containing it to drift onto them, since even minute quantities of the spray may cause severe injury during both growing and dormant periods. (Coarse sprays are less likely to drift.) Accordingly, applications by airplane, ground rigs and hand dispensers should be carried out only when there is no hazard from drift. Do not apply by airplane in the vicinity of cotton, grapes or other desirable 2,4-D Sodium Salt susceptible vegetation. Do not contaminate irrigation ditches or water used for domestic or irrigation purposes.

Do not store near fertilizers, seeds, insecticides or fungicides. Excessive amounts of 2,4-Dichlorophenoxyacetic acid in the soil may temporarily inhibit seed germination or plant growth.

To avoid injury to desirable plants, do not store, handle or apply other agricultural chemicals with the same containers or equipment used for 2,4-D Sodium Salt.

Do not reuse container. Destroy by burying in a safe place away from water supplies. Do not burn.

SOL BY
E-Z-FLO CHEMICAL COMPANY
DIVISION OF KIRSTO COMPANY
COLUMBUS, OHIO LANSING, MICHIGAN

E-Z FLO 2,4-D

CAUTION:
KEEP OUT OF THE REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

ACCEPT

January 22,

UNDER THE FEDERAL
FUNGICIDE AND RODENT
FOR ECONOMIC POISON
ED UNDER NO. 94.4% 6

ACTIVE INGREDIENT:

**2,4-Dichlorophenoxyacetic acid,
sodium salt, monohydrate***

INERT INGREDIENTS:

5.6%

100.0%

*Equivalent to 80 % 2,4-Dichlorophenoxyacetic acid.

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

DIRECTIONS FOR USE

WEED CONTROL IN ASPARAGUS: To control wild morning glory and certain other broad-leaved weeds in asparagus plantings, use 2 to 2 1/2 pounds per acre in approximately 60 gallons of water. In airplane application, use 2 pounds in approximately 12 gallons of water per acre. A wetting agent can be added if desired in both ground and airplane sprays. Apply on actively growing weeds, usually in April or May. Make no more than one application at least 1 month apart during the harvest season. Post-harvest spraying should be only by ground rig, using drop nozzles to avoid spraying the fern. For young seedlings, do not exceed 1.25 lbs per acre. Be sure that use of this product conforms to all applicable regulations.

E-Z-FLO CHEMICAL
DIVISION
COLUMBUS, OHIO