

E-Z FLO M.A.P. 2-E



Net Contents 1 Gallon
(Contains 2 lbs. methyl azinphos per gallon)



DANGER: KEEP OUT OF REACH OF CHILDREN
(See Side Panel for Additional Warnings and First Aid Instructions)

ACTIVE INGREDIENTS:

O, O-Dimethyl S- [(4-oxo-1,2,3-Benzyothiazin-3 (4H)-yl) Methyl] phosphorodithioate.....	22.7%
Xylene.....	70.6%
INERT INGREDIENTS:	6.7%
Total	100.0%

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

411-26

EPA Est. 635-MI-1; OH-1


E.P.A. Reg. No. 635-667

SOLD BY

E-Z FLO CHEMICAL COMPANY

Division of Kirsto Company

Lansing, Michigan • Columbus, Ohio



DANGER

POISON

CONTAINER DISPOSAL: DO NOT RE-USE THE CONTAINER- Completely empty the contents and bury the unused chemical at least 18 inches deep in an isolated location away from water supplies. Rinse empty drum by filling with water and add 1 cup of household lye to each 5 gallons of water. Bury rinse solution at least 18 inches deep in an isolated area away from water supply. Punch holes in the top and bottom of the can, crush the can, and bury deeply in an isolated location. Do not use container in any connection with feed, food, and drinking water.

RESTRICTIONS: Do not use on other crops used for food or forage. Use only according to label directions. Application at rates above those shown may result in illegal crop residues. Do not graze livestock in treated orchards or groves for 21 days after treatment. Do not treat food crops grown in the greenhouse. Do not apply when weather conditions favor drift from areas treated. Workers who must enter treated fields within 2 days of application should wear protective clothing. Since certain states may require specific worker re-entry periods for various crops treated, consult your state regulations or your State Department of Agriculture for information.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do

Continued on panel to right -

DIRECTIONS FOR USE (Continued)

VEGETABLES

BEANS (snap & dry) For the control of Aphids, Bean leaf beetle, Green cloverworm, Leafhoppers, Leaf miners, Stink bugs, and Velvet bean caterpillar: Apply 1 1/2 - 2 pts. per acre. For the control of Mexican bean beetle, Spotted cucumber beetle, Striped cucumber beetle, Tarnished plant bug, and Western striped cucumber beetle: Apply 2 pts. per acre. Apply specified dosage per acre by air or ground equipment in sufficient water for complete coverage but not less than 1 gallon per acre. Repeat as necessary. Do not apply to snap beans within 7 days of harvest nor to dry beans within 30 days of harvest. Do not exceed 4 applications on dry beans. Do not feed or ensile treated forage.

DIRECTIONS FOR USE

E-Z FLO M.A.P. 2-E



Net Contents 1 Gallon
(Contains 2 lbs. methyl azinphos per gallon)



POISON
DANGER: KEEP OUT OF REACH OF CHILDREN

(See Side Panel for Additional Warnings and Antidote Statements)

ACTIVE INGREDIENTS:

O, O-Dimethyl S- [(4-oxo-1,2,3-Benzothiazin-3 (4H)-yl) Methyl] phosphorodithioate.....22.7%
Xylene.....70.6%

INERT INGREDIENTS:.....6.7%
Total 100.0%

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

411-26

EPA Est. 635-MI-1; OH-1


E.P.A. Reg. No. 635-667

SOLD BY

E-Z FLO CHEMICAL COMPANY

Division of Kirsto Company

Lansing, Michigan • Columbus, Ohio



ENVIRONMENTAL CAUTIONS (continued)

not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Destroy Container When Empty. Never Re-use.

DIRECTIONS FOR USE

DANGER

Poisonous if swallowed, inhaled or absorbed through the skin. Do not get in eyes or on skin. Do not breathe fumes or spray mist. Spray operator should work to windward to stay out of drift or mist. Wear protective clothing, natural rubber gloves and goggles. A mask or respirator recommended by the U. S. Bureau of Mines should also be worn when handling the concentrate. Do not contaminate feed or food stuffs. Keep out of reach of children and domestic animals. Keep all unprotected persons out of the operating area or vicinity where there may be danger of drift. Wash hands, arms and face thoroughly with soap and warm water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

SYMPTOMS OF POISONING: A sense of "tightness" in the chest. Sweating. Contracted pupils. Stomach pains. Vomiting and diarrhea.

TREATMENT: In case of poisoning call a physician immediately. Have patient lie down and keep quiet. If swallowed, induce vomiting by giving a tablespoonful of salt in a glass of warm water. Administer milk or water freely and again induce vomiting until vomit fluid is clear. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash with flowing water for at least 15 minutes.

POISON

TO PHYSICIAN: ANTIDOTE- Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. 2-PAA is also antidotal, and may be administered in conjunction with atropine. Compound inhibits cholinesterase, resulting in stimulation of the central nervous system, the parasympathetic nervous system and the somatic motor nerves. Do not give morphine. Watch for pulmonary edema, which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically.

CONTAINER DISPOSAL: DO NOT RE-USE THE CONTAINER- Completely empty the contents and bury the unused chemical at least 18 inches deep in an isolated location away from water supplies. Rinse empty drum by filling with water and add 1 cup of household lye to each 5 gallons of water. Bury rinse solution at least 18 inches deep in an isolated area away from water supply. Punch holes in the top and bottom of the can, crush the can, and bury deeply in an isolated location. Do not use container in any connection with feed, food, and drinking water.

RESTRICTIONS: Do not use on other crops used for food or storage. Use only according to label directions. Application at rates above those shown may result in illegal crop residues. Do not graze livestock in treated orchards or groves for 21 days after treatment. Do not treat food crops grown in the greenhouse. Do not apply when weather conditions favor drift from areas treated. Workers who must enter treated fields within 2 days of application should wear protective clothing. Since certain states may require specific worker re-entry periods for various crops treated, consult your state regulations or your State Department of Agriculture for information.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do

Continued on panel to right-

DIRECTIONS FOR USE (Continued)

VEGETABLES

BEANS (snap & dry): For the control of Aphids, Bean leaf beetle, Green cloverworm, Leafhoppers, Leaf miners, Stink bugs, and Velvet bean caterpillar: Apply 1 1/2 -2 pts. per acre. For the control of Mexican bean beetle, Spotted cucumber beetle, Striped cucumber beetle, Tarnished plant bug, and Western striped cucumber beetle: Apply 2 pts. per acre. Apply specified dosage per acre by air or ground equipment in sufficient water for complete coverage but not less than 1 gallon per acre. Repeat as necessary. Do not apply to snap beans within 7 days of harvest nor to dry beans within 30 days of harvest. Do not exceed 4 applications on dry beans. Do not feed or ensile treated forage.

INERT INGREDIENTS:.....6.7%
Total 100.0%

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

411-26


EPA Est. 635-MI-1; OH-1

E.P.A. Reg. No. 635-667

SOLD BY

E-Z FLO CHEMICAL COMPANY

Division of Kirsto Company
Lansing, Michigan • Columbus, Ohio



ENVIRONMENTAL CAUTIONS (continued)

not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Destroy Container When Empty. Never Re-use.

DIRECTIONS FOR USE

This insecticide forms an emulsion when diluted with water, and is suitable for use in all power-operated ground sprayers and aircraft sprayers. To mix with water, pour the required amount into the full amount of water and then agitate.

DOSAGE: Use specified dosage in the amount of water necessary to give complete coverage of foliage. The type of equipment used will determine the concentration required.

SPRAYING: Work to windward. Protect sprayer operators from drift or mist. When low volumes of spray are applied, complete coverage and thorough application are essential for most effective results. Schedule applications in accordance with local conditions. Consult your State Agricultural Experiment Station or Extension Service for specific timing of applications in your area.

Compatible with many of the commonly used fungicides and insecticides, but is of questionable compatibility (physical) with the following: Cyrex, oils (dormant and summer) and wettable sulfur. Incompatible with Bordeaux, lime sulfur, lime, and zinc sulfate plus lime.

FRUIT

APRICOTS, PEACHES: For the control of European fruit lecanium scale, Forbes scale, Lesser peach tree borer, Mites, Peach tree borer, Plum curculio, Red-banded leaf roller, San Jose scale, Stink bug, and Tarnished plant bug: Apply 1 to 1½ pints per 100 gal. of water as a full coverage spray, using not more than 800 gals. per acre. Apply at petal fall and every 10-14 days thereafter until control is achieved. For scale control, apply when crawlers are present. Do not apply more than 8 times per season, nor within 21 days of harvest. For the control of oriental fruit moth and peach twig borer: Apply 1½ pts. per 100 gals. in not more than 800 gals. of water per acre only before mid-season. For control of peach tree borer: Apply 2 to 3 sprays to trunk from ground to scaffold limbs, timed with moth flight.

CHERRIES: For the control of Eye-spotted bud moth, Forbes scale, Fruit flies, Fruit tree leaf roller, Lesser peach tree borer, Plum curculio, and San Jose scale: Apply 1 to 2 pts. in 100 gals. of water as a full coverage spray using not more than 800 gals. of finished spray per acre. Apply at petal fall and every 10-14 days thereafter until control is achieved. For the control of Cherry Leaf miner and Mineola moth: Apply 1 pt. in 100 gals. of water. Do not apply more than 8 times per season nor within 15 days of harvest.

GRAPES: For the control of Grape berry moth, Grape mealybug, Leafhoppers, Red-banded leaf rollers, and Thrips: Apply 1-2 pts. in 100 gallons of water as a full coverage spray using not more than 300 gallons of finished spray per acre. Repeat as necessary but not more than 3 times per season. Minimum dosage specified may be applied up to harvest. Higher rates up to 1½ pints require a 10-day interval to harvest and above 1½ pints a 28-day interval to harvest.

STRAWBERRIES: For the control of Aphids, Meadow spittlebug, Oblique-banded leaf roller, Obscure root weevil, Omnivorous leaf tier, Pea leaf weevil, and Strawberry leaf rollers: Apply 2 pts. in 200 gals. of water per acre as full coverage spray. Where ground conditions dictate an air application, use specified rate in a minimum of 1 gallon of water per acre. Repeat as necessary. Do not apply within 5 days of harvest.

Continued on panel to right-

container in any connection with feed, food, and drinking water.

RESTRICTIONS: Do not use on other crops used for food or storage. Use only according to label directions. Application at rates above those shown may result in illegal crop residues. Do not graze livestock in treated orchards or groves for 21 days after treatment. Do not treat food crops grown in the greenhouse. Do not apply when weather conditions favor drift from areas treated. Workers who must enter treated fields within 2 days of application should wear protective clothing. Since certain states may require specific worker re-entry periods for various crops treated, consult your state regulations or your State Department of Agriculture for information.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do

Continued on panel to right-

DIRECTIONS FOR USE (Continued)

VEGETABLES

BEANS (snap & dry): For the control of Aphids, Bean leaf beetle, Green cloverworm, Leafhoppers, Leaf miners, Stink bugs, and Velvet bean caterpillar: Apply 1½-2 pts. per acre. For the control of Mexican bean beetle, Spotted cucumber beetle, Striped cucumber beetle, Tarnished plant bug, and Western striped cucumber beetle: Apply 2 pts. per acre. Apply specified dosage per acre by air or ground equipment in sufficient water for complete coverage but not less than 1 gallon per acre. Repeat as necessary. Do not apply to snap beans within 7 days of harvest nor to dry beans within 30 days of harvest. Do not exceed 4 applications on dry beans. Do not feed or ensile treated forage.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, and CAULIFLOWER: For the control of Aphids, Cabbage looper, Diamond-back moth, and Imported cabbageworm: Apply 2-3 pts. per acre by air or ground equipment in sufficient water for complete coverage but not less than 1 gallon per acre. Repeat as necessary. Do not apply within 7 days of harvest for brussels sprouts, 15 days of harvest for broccoli and cauliflower, nor within 21 days of harvest for cabbage. For the control of cabbage maggot: Apply ½ pint in 50 gallons of water. Apply 4 to 6 ounces of this emulsion per plant immediately after transplanting. **CELERY:** For the control of Aphids, Leaf miners, Leafhoppers, Spittlebugs, and Tarnished plant bug: Apply 2 pts in 100 gallons of water as a full coverage spray using not more than 200 gallons of finished spray per acre. Repeat as necessary but not within 14 days of harvest.

EGGPLANT: For the control of Leafminers: Apply 1½-2 pts. per acre. For the control of European corn borer and Flea beetles: Apply 2 pts. per acre. Apply specified dosage per acre by air or ground equipment in sufficient water to give complete coverage but not less than 1 gallon per acre. Repeat as necessary. Do not apply after fruit set.

MELONS (Including Honeydew melons, Muskmelons (cantaloupe), Watermelons, and other melons): For the control of Leafhoppers and Leaf miners: Apply 1½-2 pts. per acre. For the control of Rindworms, Spotted cucumber beetle, Striped cucumber beetle, and Western-striped cucumber beetle: Apply 2 pts. per acre. Apply specified dosage per acre by air or ground equipment in sufficient water to give complete coverage but not less than 1 gallon per acre. Repeat as necessary up to the day of harvest but not more than 4 times per season.

ONIONS (green & dry): For the control of thrips: Apply 2-3 pts. per acre by air or ground equipment in sufficient water to give complete coverage but not less than 1 gallon per acre. Do not apply more than 3 times per season nor within 28 days of harvest of dry onions or 7 days of harvest of green onions.

PEPPERS: For the control of leaf miners: Apply 1½-2 pts. per acre. For the control of European corn borer, and Flea beetles: Apply 2 pts. per acre. Apply specified dosage per acre by air or ground equipment in sufficient water to give complete coverage but not less than 1 gallon per acre. A maximum of 4 treatments may be made up to within 3 days of harvest. Where more than 4 applications are required, do not apply last spray within 14 days of harvest.

POTATOES: For the control of Colorado potato beetle: Apply 1½ pts. per acre. For the control of Banded cucumber beetle and Leaf miners: Apply 1½-2 pts. per acre. For the control of Aphids, European corn borer, Flea beetle, Leafhoppers, Spittlebugs, and Tarnished plant bug: Apply 2 to 3 pts. per acre. For the control of Tuberworm: Apply 2½ to 3 pts. per acre. Apply specified dosage per acre by air or ground equipment in sufficient water for complete coverage but not less than 1 gallon per acre. Repeat as necessary. Do not apply within 14 days of harvest.

TOMATOES: For the control of Colorado potato beetle: Apply 1½ pts. per acre. For the control of Banded cucumber beetle, Drosophila, Green stink bug, Leaf miners, and Whitefly: Apply 1½ to 2 pts. per acre. For the control of Aphids, European corn borer, Flea beetles, Grasshoppers, Leafhoppers, and Thrips: Apply 2 to 3 pts. per acre. For the control of Tuberworm: Apply 2½ to 3 pts. per acre. For the control of Tomato Hornworm: Apply 3 pts. per acre. For the control of Corn earworm, Fruitworm, Pinworm, and Yellow-striped armyworm: Apply 3 to 6 pts. per acre. Apply specified dosage per acre by air or ground equipment in sufficient water for complete coverage but not less than 1 gallon per acre. Repeat as necessary. The high rates should be used where heavy infestations of late instar lepidopterous larvae (large worms) and pinworms are present. Rates of 3 pints or less per acre may be applied up to day of harvest. Rates above 3 pints per acre require an interval of 14 days between application and harvest.