


E-Z-Flo Crop Guard



CAUTION: KEEP OUT OF THE REACH OF CHILDREN

Actual Ingredients: Methidathion 20.0%
 Insecticide active ingredient 4.0% (1.0% active ingredient)
 Net Weight 10.0 lbs (4.5 kg)
 Net Volume 1.0 gal (3.8 L)
 Contains 100% active ingredient

SOLD BY
E-Z FLO CHEMICAL COMPANY
 Division of Kirsto Company
 Lansing, Michigan • Columbus, Ohio

CAUTION

Avoid contact with skin, eyes and clothing. Avoid contamination of food and feed. Wash thoroughly after handling. Harmful if swallowed. May be absorbed through skin. Avoid breathing spray mist.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes and containers. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not re-use empty container. Destroy by perforating or crushing and burying in a safe place.

Store at temperatures above 32° F

GENERAL INFORMATION

This product is a systemic insecticide. It is absorbed by the plant and moves through the vascular system to the growing points of the plant. It is effective against a wide range of insects including beet beetles, leafhoppers, and grasshoppers.

Use this product only on crops listed on the label. Do not use on crops not listed. Do not use on crops if the crop is in bloom or if the crop is to be used for human consumption.

APPROXIMATE RATES

Apply 2.0 to 4.0 lbs of product per acre in 100 gallons of water. For broadcast applications, use 2.0 lbs per acre. For spot applications, use 4.0 lbs per acre. Do not apply more than 4.0 lbs per acre.

Apply this product to the soil or to the foliage of the plant. Do not apply to the fruit or to the seed. Do not apply to the soil if the soil is to be used for human consumption.

Do not apply this product to crops if the crop is in bloom or if the crop is to be used for human consumption.

DIRECTIONS FOR USE

Field Crops: Alfalfa, Clover, Cowpea, Soybean, Vetch, and other legumes. Apply 2.0 to 4.0 lbs per acre in 100 gallons of water. Do not apply to crops if the crop is in bloom or if the crop is to be used for human consumption.

DIRECTIONS FOR USE (continued)

Small Grains: For the control of the insects on the following crops, apply 2.0 to 4.0 lbs per acre in 100 gallons of water per acre as a thorough coverage spray when insects first appear. Repeat applications at 7-10 day intervals if necessary.

Vegetable Crops: Beans, Leafhoppers, Spittlebugs, and other insects. Do not apply within 14 days of harvest.

Other Crops: Blueberries, Grape Berry Moths, Grape Leaf Folders, Grape Leaf Skeletonizers, Japanese Beetles, Leafhoppers, Pacific Spruce Borer, Pear Thrips, Pear Leaf Miner, Pear Leaf Skeletonizer, Pear Leaf Rollers, Pear Leaf Tortrix, Pear Leaf Webworm, Pear Leaf Miner, Pear Leaf Skeletonizer, Pear Leaf Rollers, Pear Leaf Tortrix, Pear Leaf Webworm.

Home Gardens: For the control of insects on the following crops, apply 2.0 to 4.0 lbs per acre in 100 gallons of water per acre as a thorough coverage spray when insects first appear. Repeat applications at 7-10 day intervals if necessary. Do not apply within 14 days of harvest.

Lawns: For the control of insects on the following crops, apply 2.0 to 4.0 lbs per acre in 100 gallons of water per acre as a thorough coverage spray when insects first appear. Repeat applications at 7-10 day intervals if necessary. Do not apply within 14 days of harvest.

Vegetable Crops: For the control of the insects on the following crops, use 2.0 to 4.0 lbs per acre in sufficient water for thorough coverage. Begin treatment at first sign of insect infestation and repeat applications at 7-10 day intervals if necessary. Do not apply to crops if the crop is in bloom or if the crop is to be used for human consumption.

DIRECTIONS FOR USE (continued)

Other Crops: Apples, Apricots, Blackberries, Blueberries, Cherries, Citrus, Grapes, Peaches, Pears, Plums, Raspberries, Strawberries, and other fruits. Do not apply to crops if the crop is in bloom or if the crop is to be used for human consumption.

Other Crops: Apples, Apricots, Blackberries, Blueberries, Cherries, Citrus, Grapes, Peaches, Pears, Plums, Raspberries, Strawberries, and other fruits. Do not apply to crops if the crop is in bloom or if the crop is to be used for human consumption.

Other Crops: Apples, Apricots, Blackberries, Blueberries, Cherries, Citrus, Grapes, Peaches, Pears, Plums, Raspberries, Strawberries, and other fruits. Do not apply to crops if the crop is in bloom or if the crop is to be used for human consumption.

Other Crops: Apples, Apricots, Blackberries, Blueberries, Cherries, Citrus, Grapes, Peaches, Pears, Plums, Raspberries, Strawberries, and other fruits. Do not apply to crops if the crop is in bloom or if the crop is to be used for human consumption.

Other Crops: Apples, Apricots, Blackberries, Blueberries, Cherries, Citrus, Grapes, Peaches, Pears, Plums, Raspberries, Strawberries, and other fruits. Do not apply to crops if the crop is in bloom or if the crop is to be used for human consumption.

Other Crops: Apples, Apricots, Blackberries, Blueberries, Cherries, Citrus, Grapes, Peaches, Pears, Plums, Raspberries, Strawberries, and other fruits. Do not apply to crops if the crop is in bloom or if the crop is to be used for human consumption.

Other Crops: Apples, Apricots, Blackberries, Blueberries, Cherries, Citrus, Grapes, Peaches, Pears, Plums, Raspberries, Strawberries, and other fruits. Do not apply to crops if the crop is in bloom or if the crop is to be used for human consumption.

Other Crops: Apples, Apricots, Blackberries, Blueberries, Cherries, Citrus, Grapes, Peaches, Pears, Plums, Raspberries, Strawberries, and other fruits. Do not apply to crops if the crop is in bloom or if the crop is to be used for human consumption.

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

SOLD BY
E-Z FLO CHEMICAL COMPANY
Division of Kirsto Company
Lansing, Michigan • Columbus, Ohio

GENERAL INFORMATION

This product is a highly toxic insecticide. It is effective against a wide range of insects, including beetles, flies, and mosquitoes. It is used for the control of insects in various agricultural and domestic settings. The product is available in a spray form and should be applied according to the directions on the label. It is important to use this product carefully to avoid contamination of water and other resources.

DIRECTIONS FOR USE

Apply this product to the insects to be controlled. Use a spray nozzle to apply the product to the insects. The product should be applied to the insects directly, and not to the surrounding area. The product is effective against a wide range of insects, including beetles, flies, and mosquitoes. It is important to use this product carefully to avoid contamination of water and other resources.

DIRECTIONS FOR USE

For the control of the insects in the following crops, apply 2 1/2 to 5 1/2 gallons of water per acre as a thorough spray. Repeat applications at 7-10 day intervals, if necessary.

For the control of the insects in the following crops, apply 1/2 to 1 1/2 gallons of water per acre as a thorough spray. Repeat applications at 7-10 day intervals, if necessary.

For the control of the insects in the following crops, apply 1/2 to 1 1/2 gallons of water per acre as a thorough spray. Repeat applications at 7-10 day intervals, if necessary.

For the control of the insects in the following crops, apply 1/2 to 1 1/2 gallons of water per acre as a thorough spray. Repeat applications at 7-10 day intervals, if necessary.

from areas treated. Do not apply where runoff is likely to occur. Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes and containers. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

This product is highly toxic. If bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not use empty container. Destroy by perforating, crushing and burying in a safe place.

Store at temperatures above 32° F

DIRECTIONS FOR USE

Control of Aphids, Fleas, Flies, Mosquitoes, Beetles, House Flies, and other insects. Do not apply with water.

Control of Beetles, Japanese Beetles, Leafhoppers, and other insects. Do not apply with water.

Control of Beetles, Japanese Beetles, Leafhoppers, and other insects. Do not apply with water.

Control of Beetles, Japanese Beetles, Leafhoppers, and other insects. Do not apply with water.