



E-Z FLO DIAZINON 14 G INSECTICIDE

CAUTION:
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT
O, O-diethyl O-(12-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 14.3%
INERT INGREDIENTS 85.7%
100.0%
Diazinon® trademark of CIBA-GEIGY

CAUTION

Harmful if swallowed. Avoid contamination of food and feed. Do not store near food or feed products. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. May be absorbed through skin.

This product is toxic to fish, shrimp, crabs, birds and other wildlife. Birds feeding in treated areas may be killed. Keep out of lakes, streams, ponds, wet marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as directed on this label.

Do not reuse container. Destroy it by puncturing or burning. Avoid reuse of container.

51-29 EPA Est. No. 149-100001-1 P. A. Reg. No. 149-100001-1

DIRECTIONS FOR USE

Applications to the soil should be worked into the top few inches of soil by plowing, disking, or other means. Do not apply to water. Do not apply to water.

FIELD AND PASTURE USE

BEANS—Pinto, Lesser Green, and Black Beans. Apply 14-28 lbs. per acre as broadcast soil treatment 1-2 weeks before planting.

CORN—Corn Rootworm, European Corn Borer, and European Cornborer. Apply 14-28 lbs. per acre as broadcast soil treatment 1-2 weeks before planting.

CORN—Corn Rootworm, European Corn Borer, and European Cornborer. Apply 14-28 lbs. per acre as broadcast soil treatment 1-2 weeks before planting.

CORN—Corn Rootworm, European Corn Borer, and European Cornborer. Apply 14-28 lbs. per acre as broadcast soil treatment 1-2 weeks before planting.

CORN—Corn Rootworm, European Corn Borer, and European Cornborer. Apply 14-28 lbs. per acre as broadcast soil treatment 1-2 weeks before planting.

CORN—Corn Rootworm, European Corn Borer, and European Cornborer. Apply 14-28 lbs. per acre as broadcast soil treatment 1-2 weeks before planting.

CORN—Corn Rootworm, European Corn Borer, and European Cornborer. Apply 14-28 lbs. per acre as broadcast soil treatment 1-2 weeks before planting.

CORN—Corn Rootworm, European Corn Borer, and European Cornborer. Apply 14-28 lbs. per acre as broadcast soil treatment 1-2 weeks before planting.

plants to have granules falling to the ground by the end of 2-3 inches of soil into the corn row with a disc harrow or cultivator. Repeat application in corn when 7 days later if needed. For control of European cornborer, to control second-generation corn borers apply 7-14 lbs. per acre as a level when field corn is 100 egg masses per 100 plants. Fall Armyworms. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when armyworms first appear. Repeat application if necessary. Southwestern Corn Borers. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when second-generation corn borers first appear. Repeat application if necessary.

NOTE—Do not feed treated forage to dairy or beef cattle or sheep for 30 days following application. Corn may be grazed immediately. Temporary spitting of foliage sometimes occurs following application.

CUTWORMS—Broadcast 14-28 lbs. per acre. For surface cutworms apply just prior to planting or when larvae first appear and work into soil 1-2 inches. Do not apply after layby. For subterranean cutworms apply just prior to planting and work into soil 4-6 inches. Lesser Greenstalk Borers. Apply 8-17 oz. per 1,000 linear feet of row in a 10-inch band centered over the row. Make application at planting time when crop begins to emerge. Cover the treated band with 2 inches of soil. Apply amount per acre recommended according to row spacing chart for basal treatment. Seed Corn Maggots. Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2 inches. Garden Symphyla and Beetles. Broadcast 7 lbs. per acre just prior to planting and work into soil 4-6 inches. This preplant treatment will also control other soil insects such as cutworms, surface and subterranean and wireworms. Wireworms. Broadcast 27-28 lbs. per acre before planting. Work into soil 4-6 inches.

SUGGESTED RATES AND ROW SPACING

Row Spacing	Linear Feet of Row	Amount of Diazinon 14 G per acre
40"	1,497	36-60
42"	1,429	37-66
44"	1,364	38-72
46"	1,300	39-78
48"	1,237	40-84
50"	1,175	41-90
52"	1,114	42-96
54"	1,053	43-102

NOTE—For broadcast applications, the amount of Diazinon 14 G per acre should be adjusted to the row spacing used.

FIELD USE—Broadcast 14-28 lbs. per acre over the soil 1-2 weeks before planting. For control of European cornborer, to control second-generation corn borers apply 7-14 lbs. per acre as a level when field corn is 100 egg masses per 100 plants. Fall Armyworms. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when armyworms first appear. Repeat application if necessary. Southwestern Corn Borers. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when second-generation corn borers first appear. Repeat application if necessary.

FIELD USE—Broadcast 14-28 lbs. per acre over the soil 1-2 weeks before planting. For control of European cornborer, to control second-generation corn borers apply 7-14 lbs. per acre as a level when field corn is 100 egg masses per 100 plants. Fall Armyworms. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when armyworms first appear. Repeat application if necessary. Southwestern Corn Borers. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when second-generation corn borers first appear. Repeat application if necessary.

FIELD USE—Broadcast 14-28 lbs. per acre over the soil 1-2 weeks before planting. For control of European cornborer, to control second-generation corn borers apply 7-14 lbs. per acre as a level when field corn is 100 egg masses per 100 plants. Fall Armyworms. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when armyworms first appear. Repeat application if necessary. Southwestern Corn Borers. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when second-generation corn borers first appear. Repeat application if necessary.

FIELD USE—Broadcast 14-28 lbs. per acre over the soil 1-2 weeks before planting. For control of European cornborer, to control second-generation corn borers apply 7-14 lbs. per acre as a level when field corn is 100 egg masses per 100 plants. Fall Armyworms. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when armyworms first appear. Repeat application if necessary. Southwestern Corn Borers. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when second-generation corn borers first appear. Repeat application if necessary.

FIELD USE—Broadcast 14-28 lbs. per acre over the soil 1-2 weeks before planting. For control of European cornborer, to control second-generation corn borers apply 7-14 lbs. per acre as a level when field corn is 100 egg masses per 100 plants. Fall Armyworms. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when armyworms first appear. Repeat application if necessary. Southwestern Corn Borers. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when second-generation corn borers first appear. Repeat application if necessary.

FIELD USE—Broadcast 14-28 lbs. per acre over the soil 1-2 weeks before planting. For control of European cornborer, to control second-generation corn borers apply 7-14 lbs. per acre as a level when field corn is 100 egg masses per 100 plants. Fall Armyworms. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when armyworms first appear. Repeat application if necessary. Southwestern Corn Borers. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when second-generation corn borers first appear. Repeat application if necessary.

FIELD USE—Broadcast 14-28 lbs. per acre over the soil 1-2 weeks before planting. For control of European cornborer, to control second-generation corn borers apply 7-14 lbs. per acre as a level when field corn is 100 egg masses per 100 plants. Fall Armyworms. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when armyworms first appear. Repeat application if necessary. Southwestern Corn Borers. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when second-generation corn borers first appear. Repeat application if necessary.

FIELD USE—Broadcast 14-28 lbs. per acre over the soil 1-2 weeks before planting. For control of European cornborer, to control second-generation corn borers apply 7-14 lbs. per acre as a level when field corn is 100 egg masses per 100 plants. Fall Armyworms. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when armyworms first appear. Repeat application if necessary. Southwestern Corn Borers. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when second-generation corn borers first appear. Repeat application if necessary.

FIELD USE—Broadcast 14-28 lbs. per acre over the soil 1-2 weeks before planting. For control of European cornborer, to control second-generation corn borers apply 7-14 lbs. per acre as a level when field corn is 100 egg masses per 100 plants. Fall Armyworms. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when armyworms first appear. Repeat application if necessary. Southwestern Corn Borers. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when second-generation corn borers first appear. Repeat application if necessary.

FIELD USE—Broadcast 14-28 lbs. per acre over the soil 1-2 weeks before planting. For control of European cornborer, to control second-generation corn borers apply 7-14 lbs. per acre as a level when field corn is 100 egg masses per 100 plants. Fall Armyworms. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when armyworms first appear. Repeat application if necessary. Southwestern Corn Borers. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when second-generation corn borers first appear. Repeat application if necessary.

FIELD USE—Broadcast 14-28 lbs. per acre over the soil 1-2 weeks before planting. For control of European cornborer, to control second-generation corn borers apply 7-14 lbs. per acre as a level when field corn is 100 egg masses per 100 plants. Fall Armyworms. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when armyworms first appear. Repeat application if necessary. Southwestern Corn Borers. Apply 7-14 lbs. per acre over the corn as granules fall into whorls when second-generation corn borers first appear. Repeat application if necessary.

FORAGE CROPS—Alfalfa, Lucerne, Clover, and Lespedeza. C. worms. Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms. 3-6 inches to control subterranean cutworms.

VEGETABLE CROP INSECTS

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, AND CAULIFLOWER—Root Maggots. Broadcast 14-21 lbs. per acre before planting or transplanting. Work into soil 3-4 inches.

ONIONS—Onion Maggots. Apply 14-28 lbs. per acre as broadcast soil treatment just prior to planting. Thoroughly mix the chemical into the top 3-4 inches of soil by disking, harrowing or other suitable means.

RADISHES—Root Maggots. Apply 10-14 lbs. per acre (21-28 lbs. per 1,000 linear feet of row) at the seed furrow at planting time.

SWEET CORN—See Corn under Field and Forage Crops.
TOMATOES—Tomato Pinworm, Drosophila spp. Broadcast 7 lbs. per acre when flies first appear. Repeat at 10 day intervals if as required.

OTHER VEGETABLE INSECTS

Crop	Insect	Rate
Broccoli	Root Maggots	14-21 lbs. per acre
Brussels Sprouts	Root Maggots	14-21 lbs. per acre
Cabbage	Root Maggots	14-21 lbs. per acre
Cauliflower	Root Maggots	14-21 lbs. per acre
Lettuce	Root Maggots	14-21 lbs. per acre
Lima Beans	Root Maggots	14-21 lbs. per acre
Muskmelon	Root Maggots	14-21 lbs. per acre
Peas	Root Maggots	14-21 lbs. per acre
Peppers	Root Maggots	14-21 lbs. per acre
Pine Beans	Root Maggots	14-21 lbs. per acre
Snap Beans	Root Maggots	14-21 lbs. per acre
Swiss Chard	Root Maggots	14-21 lbs. per acre
Tomatoes	Tomato Pinworm	7 lbs. per acre
Winter Squash	Root Maggots	14-21 lbs. per acre

Broadcast 14 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms. 3-6 inches to control subterranean cutworms. Potatoe. Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 4-6 inches.

CARDEN SYMPHYLID INSECTS

Crop	Insect	Rate
Cauliflower	Root Maggots	14-21 lbs. per acre
Corn	Root Maggots	14-21 lbs. per acre
Peas	Root Maggots	14-21 lbs. per acre
Pine Beans	Root Maggots	14-21 lbs. per acre
Radishes	Root Maggots	10-14 lbs. per acre
Tomatoes	Tomato Pinworm	7 lbs. per acre

Broadcast 14 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms. 3-6 inches to control subterranean cutworms. Potatoe. Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 4-6 inches.

Broadcast 14 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms. 3-6 inches to control subterranean cutworms. Potatoe. Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 4-6 inches.

Broadcast 14 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms. 3-6 inches to control subterranean cutworms. Potatoe. Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 4-6 inches.

Broadcast 14 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms. 3-6 inches to control subterranean cutworms. Potatoe. Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 4-6 inches.

Broadcast 14 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms. 3-6 inches to control subterranean cutworms. Potatoe. Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 4-6 inches.

Broadcast 14 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms. 3-6 inches to control subterranean cutworms. Potatoe. Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 4-6 inches.

Broadcast 14 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms. 3-6 inches to control subterranean cutworms. Potatoe. Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 4-6 inches.

Broadcast 14 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms. 3-6 inches to control subterranean cutworms. Potatoe. Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 4-6 inches.

Broadcast 14 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms. 3-6 inches to control subterranean cutworms. Potatoe. Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 4-6 inches.

Broadcast 14 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms. 3-6 inches to control subterranean cutworms. Potatoe. Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 4-6 inches.

Broadcast 14 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms. 3-6 inches to control subterranean cutworms. Potatoe. Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 4-6 inches.

Broadcast 14 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms. 3-6 inches to control subterranean cutworms. Potatoe. Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 4-6 inches.

Broadcast 14 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms. 3-6 inches to control subterranean cutworms. Potatoe. Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 4-6 inches.