

### CRABGRASS CONTROL


**CRABGRASS CONTROL: TIMING IS IMPORTANT:** To prevent crabgrass, apply Chlordane 73% before seeds germinate. IN MOST AREAS IT MUST BE APPLIED BEFORE APRIL 1ST. It also has long lasting residual action in the soil. A single application will usually provide protection for the entire growing season. Do not use on newly seeded lawns or turf.

Chlordane 73%, used as directed, is harmless to established lawns. Chlordane emulsions are readily applied by means of a pressure sprayer, jar and garden hose attachment, or sprinkling can. Apply 1 1/2 pints in sufficient water (3 to 10 gallons) to drench uniformly 1,000 sq. ft. of lawn or turf. Immediately after application, wash Chlordane off grass onto soil surface. Do not allow children or pets on treated area until Chlordane has been watered into the turf and the treated area has dried completely.

**WARNING:** May be fatal if swallowed! Avoid inhaling spray. Avoid contact with eyes or clothing. In case of contact with skin, wash with soap and water.

**CAUTION:** Use ONLY at dilutions as directed. Do not contaminate any body of water nor apply to any area not specified on label. Toxic to fish and wildlife. Birds feeding in treated areas may be harmed. Do not use on animals, foods, feed crops, or in dairy barns.

**NON-WARRANTY:** Seller makes no warranty of any kind, expressed or implied, concerning the use of this product. Buyer assumes all risk in use or handling, whether in accordance with directions or not.



# CHLORDANE 73%

**ACTIVE INGREDIENTS:**  
Technical Chlordane\* 73%  
Petroleum Distillate 17%  
**INERT INGREDIENT** 10%  
\*Equivalent to 43.8% octachloro-4,7-methano-tetrahydroindane and 29.2% related compounds.

**WARNING: Keep out of reach of children.**  
See back panel for additional cautions.

### CHLORDANE 73%

CHLORDANE 73% forms an emulsion when mixed with water which facilitates spray application to large or small areas.

**FOR LAWN INSECTS:** White Grubs, Japanese Beetle Larvae, Asiatic Garden Beetle Larvae, Sod Webworms, Lawn Moths, Chinch Bugs, Wireworms, Cutworms, Chiggers, Ticks, Armyworms, and Ants in lawns. As an insecticide, dilute 2 to 4 fl. ozs. (4 to 8 tablespoons) with 5 gals. of water to treat 1,000 sq. ft. of lawn. For sandy soil, use 2 fl. ozs. (4 tablespoons); for solid type soil, use 4 fl. ozs. (1/2 cup). It is important to water the lawn thoroughly after applying Chlordane. Note **CAUTIONS** concerning children, pets and birds.

**TERMITES:** Mix 2 pints of Chlordane 73% into 25 gals. of water. Apply this emulsion at the dosage of 1 gal. per linear foot to a trench 1 to 2 feet deep and 8 to 12 inches wide dug next to the foundation. Near deep foundations, trench should be 30 inches deep, but not below the level of the footing. Apply directly to infested wooden structures. Inspect area yearly to determine possible reinfestation.

**ROACHES, SILVERFISH, SPIDERS, ANTS, IN AND AROUND DWELLINGS:** Dilute 1 1/2 fl. ozs. (3 Tablespoons) with 2 qts. of water. Thoroughly spray infested cracks and other hiding places and adjacent exposed surfaces where these pests may occur. Spot treat only. Do not contaminate food, drinking water or utensils. Repeat as necessary.

USDA Reg. No. 64215

ACCEPTED

UNDER THE FEDERAL INSECTICIDE ACT  
FOR EXPORT AND FOREIGN RESALE  
EQUIPMENT NO. 17155