

LABEL FOR:

Date sent in: 1

USDA Reg. No. 642-17

THOMPSON CHEMICALS CORPORATION  
23529 S. Figueroa St.  
Wilmington, California 90744

**FLIES AROUND DWELLINGS, Outbuildings, Picnic Areas, Animal Quarters:** 5 tbsps. to 1 gal. water. 3 to 5 tbsps. cane or dark sugar or molasses may increase effectiveness. Spray around house foundations, under porches, along fences, shrubbery and other infested areas. Use 1 gal. diluted spray to 100 sq. ft. area.

**REDUCTION OF FLY POPULATION:** In and around Garbage Cans — Paint inside of garbage can lid twice monthly with undiluted material. Use small paint brush. Paint around top of can, inside above garbage and under can.

**CAUTION.** Harmful if swallowed, inhaled or absorbed through skin. Avoid inhalation. Do not get in eyes, on skin or clothing. Wash thoroughly with soap and water after use. Change clothing immediately and wash before re-use. Keep from children and domestic animals. Malathion, upon repeated or prolonged use, may cause cholinesterase inhibition. If accidental poisoning occurs, get prompt medical aid. Atropine is antidotal.

**USE CAUTIONS.** Do not contaminate feed and foodstuffs, milk, milking equipment or drinking water. Do not use on food crops within 14 days of harvest. Injury may occur when the product is applied to Ribier grapes and to Bosc pears in the Northeast. It may also injure McIntosh and Cortland varieties of apples, if applied within four weeks of harvest. Injury may also occur when the product is applied to certain ferns, including Boston, Maidenhair and Pteris, as well as to some species of Crassula. Destroy containers completely when empty. Do not re-use.

This product is toxic to fish. Keep out of any body of water.


**NON-WARRANTY.** Seller makes no warranty, expressed or implied, concerning the use of this product. Buyer assumes all risk in use or handling, whether in accordance with directions or not.

CODE 43130



1 PINT

# MALATHION 50%



ACTIVE INGREDIENTS:	.....	By Wt.
Malathion*	.....	50%
Aromatic Petroleum Derivative Solvent	.....	39%
INERT INGREDIENT:	.....	11%
*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate		

**CAUTION: Keep out of reach of children.**  
See back panel for additional cautions.

LOS ANGELES THOMPSON CHEMICALS CORPORATION ST. LOUIS

### DIRECTIONS

Spray thoroughly covering both upper and lower leaf surfaces or other infested plant parts. Repeat as necessary.

As an aid in making other amounts of spray solution than indicated below, calculate on the basis that 3 tsps. equal 1 tbsp. and 2 tbsps. equal 1 ounce.

**ORNAMENTALS (Roses, Camellias, Azaleas, Stocks, Chrysanthemums, Evergreens and many other Flowers and Shrubs) — Aphids:** 1½ to 2 tsps. to 1 gal. water. **Red Spider Mites, Mealybugs, Woolly Aphis, White Fly, Thrips, Tarnished Plant Bug, Four-lined Leaf Bug, Rose Leafhopper, Japanese Beetle Adults:** 1 tbsp. per gal. water. **Scale Crawlers:** 1½ to 2 tbsps. per gal. water.

**FRUITS (Apples, Pears, Grapes) — Aphids:** 1½ to 2 tsps. to 1 gal. water. **Red Spider Mites, Mealybugs, Woolly Aphis, White Fly, Thrips, Tarnished Plant Bug, Four-lined Leaf Bug, Grape Leafhopper (California and Southwest only), Pear Psyllid, Japanese Beetle Adults:** 1 tbsp. per gal. water. **Scale Crawlers:** 1½ to 2 tbsps. per gal. water.

**VEGETABLES (Cole Crops, Beans, Peas, Potatoes) — Aphids:** 1½ to 2 tsps. to 1 gal. water. **Red Spider Mites, Mealybugs, White Fly, Thrips, Tarnished Plant Bug, Four-lined Leaf Bug, Bean Leafhopper, Potato Leafhopper, Japanese Beetle Adults:** 1 tbsp. per gal. water.

**CLOVER MITES (Brown Mite, Clover Spider Mite or Brown Almond Mite) in Lawns and Around Dwellings:** 2 tsps. per 1 gal. water. Spray lower foundation of house as well as ground, lawns and plants in an area 10 ft. wide along side of house. Repeat as necessary. **DO NOT USE INSIDE DWELLINGS.**

MADE IN U.S.A.

AFG

USDA Reg. No. 642-87

ACCEPTED WITH COMMENTS

*Apr 29 1967*  
*with comments*