The wike the send the not spray to the manner tall plant to the send the send this continue that plant to the send the send this to the send the plants.

to the same foodstuffs

contain port with fertilizers.
 contain or seeds. Do not use
 contain open flame

e de la compation decine de La transfer purposes

and the second section of the second

emplace tom per

PWINIY I A TO MES THE ABITATION OF THE MET THE

The New Alltha



TOMATO FIX

COURTRATE

A HORMONE-LIKE BLOOM-SET

ACTIVE INGREDIENT:

Para-Chlorophenoxyacetic Acid
as the Diethandiam de Sait 2.631

INERT INGREDIENTS 97.921

Figurvalent to Para Chlorophenoxyacetic Acid. 1.33

CAUTION: Keep out of reach of children
See back panel for additional cautions.

I GALLUN

THOMPSON CHEMICALS
DIV. of Wm. T. Thompson Co.
ST. LOUIS, MISSOURI 63132

wangley of party

DIRECTIONS

23529 S. Figueroa St. Wilmington, Calif. 90,44

CAUTION: Do not use Thompson TOMATO FIX on tomatoes grown for seed. Do not spray flower beds, vagetable, or ornamental plantings. Avoid drift of spray or vapors as this product may injure nearby susceptible plants.

Avoid contamination of feed and foodstuffs.

On not store near or transport with fertilizers, acticides, fungicides or seeds. Do not use or store near heat or open flame.

Avoid contamination of irrigation ditches or water used for domestic purposes.

Avoid contact with skin, eyes, and clothing. Wash hands after use. Keep away from pets.

Wash containers thoroughly as soon as emptied and destroy.


NON WARRANTY: Seller makes no warranty of any kind, expressed or implied, concerning the use of this product. Buyer assumes all risk in use or handling, whether in accordance with directions or not.

'ADE IN U.S.A. USDA Reg. No. 642-183

AFF

FAR 24,1911

UNDER THE PERIPAR INTERFERENCE BOUNDER HO. 117,102



Thompson TOMATO FIX aids in the production of larger tomato crops and earlier crop harvest.

DIRECTIONS

Use 1 fl. oz. (2 tbsp.) per 2 gallons of water. This makes a solution of 50 ppm of Parachlorophenoxyacetic Acid. App!y as a directed fine spray to blossom clusters when the majority of flowers in the cluster are open. Do not apply directly to leaves or fruit.

Do not spray individual clusters more than once. Do not spray leaves or fruit. Do not make more than five applications per season. Spray clusters very lightly for best results.