

## DIRECTIONS

Mix thoroughly one part of Rax Powder with 19 parts of a suitable bait material which is available, such as corn meal, rolled oats, dog food, laying mash, stock food, etc. A small amount of a bland oil, such as mineral, corn, peanut, or any other edible oil, may be used, up to 5%, when the bait is so dry that it might have a tendency to dust.

When perishable baits, such as ground meat, fish, vegetables or similar material are used, bait placements should be made in late afternoon and replaced daily with fresh bait.

All baits should be placed in locations not accessible to children, pets and domestic animals or in tamperproof bait boxes.

## RATS


Baits should be placed in amounts of from 4 oz. up to 1 lb. in locations frequented by rats or where rats have been known to feed. Baits must be protected from dogs, cats, poultry, livestock, wildlife etc., by bait boxes, cages or other suitable means.

Baits should be examined regularly and replaced when completely eaten. If bait moulds, decomposes or becomes sour it should be replaced.

An uninterrupted supply of bait should be maintained for at least 15 days.

## MICE

For controlling mice follow the same general directions as for rats except that the amount of bait placements should be tablespoon amounts (1/4 to 1/2 ounce) to be placed at 8-12 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days.



## ACTIVE INGREDIENT:

WARFARIN 3-(a-acetonylbenzyl)-4-hydroxycoumarin\* 0.5%

INERT INGREDIENTS \_\_\_\_\_ 99.5%

Total \_\_\_\_\_ 100.0%

\* Wisconsin Alumni Research Foundation Compound No. 42

FOLLOW DIRECTIONS CAREFULLY

ACCEPTED KEEP OUT OF REACH OF CHILDREN

JUN 16, 1971 CAUTION

655-19  
UNDER THE FEDERAL INSECTICIDE AND FUNGICIDE ACT FOR ECONOMIC POWDER REGISTERED UNDER THE SUBJECT TO ATTACHED CONDITIONS

DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURYING WITH WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES.

CONTENTS LBS.

## IMPORTANT

Baiting should continue as long as evidence of feeding is observed. Except in extremely heavy infestations this should be from five days to two weeks. Where a continuous source of infestation is present from nearby dumps or fields, permanent bait stations should be used and bait placed in them as often as necessary.

## CAUTION

Rax Powder contains as its active ingredient an anti-coagulant chemical therefore baits prepared with Rax Powder must be protected from or placed where humans, livestock, wildlife, poultry or pets will not eat them. In case baits are accidentally eaten, give a teaspoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

NOTE FOR PHYSICIANS: When a human has been known to have accidentally ingested any RAX POWDER, blood transfusions combined with intravenous injections and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdoses of Dicumarol†.

† Trade Mark

USDA Reg. No. 655-19



Prentiss Drug & Chemical Co., Inc.

101 WEST 31ST STREET, NEW YORK 1, N. Y.

9 SO. CLINTON STREET, CHICAGO, ILL.

655-19

06/16/1971

1/1