



prentox[®]

38

50% DDVP EMULSIFIABLE CONCENTRATE

AN INSECTICIDE FOR
FORMULATING USE ONLY.

BEST DOCUMENT AVAILABLE

ACTIVE INGREDIENTS:	
2,2-Dichlorovinyl dimethyl phosphate	46.5%
Related Compounds	3.5%
Xylene Range Aromatic	45.0%
INERT INGREDIENTS:	5.0%
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN



POISON DANGER



E.P.A. REG. NO. 655-196

E.P.A. EST. NO. 655-NJ-1

ACCEPTED

LOT # _____

NET CONTENTS _____

655-196

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

Poisonous if swallowed, inhaled, or absorbed in or eyes. Do not get in eyes or on skin. Do not breathe fumes. Rapidly absorbed through skin, wear rubber gloves, waterproof protective clothing, and boots when handling. Wear a pesticide respirator approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and the National Institute for Occupational Safety and Health for the provisions of 30 CFR Part 11.

Environmental Hazards

Product is toxic to fish, birds and other wildlife. Do not discharge into lakes, streams, ponds, or public waters in accordance with an NPDES permit. For more information contact your Regional Office of the Environmental Protection Agency.

Physical or Chemical Hazards

Store near heat or open flame.

Instructions for Formulation

Follow instructions of federal law to use this product in a manner consistent with its labeling.

Product may be used only for the formulation of emulsions. Formulators using this product are required to obtain EPA registration for their products. Refer to Technical Bulletin for more information on formulating and other information.

Note to Physician

Symptoms include weakness, headache, dizziness, blurred vision, non-reactive pin-point pupils, sweating, nausea, vomiting, diarrhea, muscle cramps.

Atropine is the specific therapeutic choice against parasympathetic nervous system. If there are signs of parasympathetic system activity, atropine should be injected at ten-minute intervals of 1 to 2 milligrams, until complete atropinization has occurred. Morphine is contraindicated.

Clear chest by postural drainage. Oxygenation may be necessary. 2-PAM is also contraindicated. Do not administer in conjunction with other patient continuously for 48 hours. Do not give if patient is on cholinesterase inhibitors. Do not give if patient is on cholinesterase inhibitors. Do not give until cholinesterase regeneration has been determined by blood test.

STATEMENT OF PRACTICAL TREATMENT

If swallowed — Call a physician immediately. Gastric lavage is indicated if material was taken internally. DO NOT INDUCE vomiting unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled — Move victim to fresh air. Apply respiration if indicated. Call a physician immediately.

If on skin — Remove contaminated clothing and wash immediately with soap and water. Call a physician immediately.

If in eyes — Flush eyes for at least 15 minutes with water. Call a physician immediately.

STORAGE AND DISPOSAL

PROHIBITIONS

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL. OPEN DUMPING IS PROHIBITED.

PESTICIDE DISPOSAL

PESTICIDE, SPRAY MIXTURE, OR RINSATE THAT CANNOT BE USED OR CHEMICALLY PROCESSED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

CONTAINER DISPOSAL

(a) RESEAL CONTAINER AND OFFER FOR RECONDITIONING, OR (b) TRIPLE RINSE (OR EQUIVALENT) AND OFFER FOR RECYCLING, RECONDITIONING, OR DISPOSAL IN APPROVED LANDFILL, OR BURY IN A SAFE PLACE IF LESS THAN 30 GALLONS; TRIPLE RINSE (OR EQUIVALENT), AND OFFER FOR RECYCLING, RECONDITIONING, OR DISPOSAL IN APPROVED LANDFILL OR BURY IN A SAFE PLACE.

GENERAL

CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES.



Manufactured by

prentiss

Drug & Chemical Co., Inc.

Plant: 338 Wilson Avenue
Newark, N.J. 07105

Office: 37th Avenue
New York, N.Y. 10001