

**JUL 5 1970**  
 FEDERAL INSECTICIDE AND RODENTICIDE BOARD  
 ECONOMIC POISON REGISTER  
 SUBJECT

**DIRECTIONS**

Mix thoroughly one part of Rax Powder with 19 parts of a suitable bait material which is available, such as corn meal, rolled oats, dog food, laying mash, stock food, etc. A small amount of a bland oil, such as mineral, corn, peanut, or any other edible oil, may be used, up to 5%, when the bait is so dry that it might have a tendency to dust.

When perishable baits, such as ground meat, fish, vegetables or similar material are used, bait placements should be made in late afternoon and replaced daily with fresh bait.

All baits should be placed in locations not accessible to children, pets and domestic animals or in tamperproof bait boxes.

**RATS**

Baits should be placed in amounts of from 4 oz. up to 1 lb. in locations frequented by rats or where rats have been known to feed. Baits must be protected from dogs, cats, poultry, livestock, etc., by bait boxes, cages or other suitable means.

Baits should be examined regularly and replaced when completely eaten. If bait moulds, decomposes or becomes sour it should be replaced.

An uninterrupted supply of bait should be maintained for at least 10 days.

**MICE**

For controlling mice follow the same general directions as for rats except that the amount of bait placements should be tablespoon amounts (1/4 to 1/2 ounce) to be placed at 8-12 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days.



**PRENTISS  
 WARFARIN  
 CONCENTRATE  
 RAX POWDER**



**IMPORTANT**

Baiting should continue as long as evidence of feeding is observed. Except in extremely heavy infestations this should be from five days to two weeks. Where a continuous source of infestation is present from nearby dumps or fields, permanent bait stations should be used and bait placed in them as often as necessary.

**CAUTION**

Rax Powder contains as its active ingredient an anti-coagulant chemical therefore baits prepared with Rax Powder must be protected from or placed where humans, livestock, poultry or pets will not accidentally eat them. In case baits are accidentally eaten, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

**NOTE FOR PHYSICIANS:** When a human has been known to have accidentally ingested any RAX POWDER, blood transfusions combined with intravenous injections and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdoses of Dicumarol†.

† Trade Mark

USDA Reg. No. 655-288

**WITH MALATHION  
 FOR MANUFACTURING PURPOSES ONLY  
 NOT FOR RESALE UNDER THIS LABEL**

**ACTIVE INGREDIENT:**

**WARFARIN 3-( $\alpha$ -acetylbenzyl)-4-hydroxycoumarin\* 0.5%**

**MALATHION (O, O-Dimethyl Dithiophosphate of Diethyl Mercaptosuccinate.) 0.2%**


**INERT INGREDIENTS 99.3%**

**Total 100%**

\* Wisconsin Alumni Research Foundation Compound No. 42

**FOLLOW DIRECTIONS CAREFULLY  
 KEEP OUT OF REACH OF CHILDREN**

**CONTENTS LBS.**



**Prentiss Manufacturing & Chemical Co. Inc.**