

A PRENTISS PEST-TESTED PRODUCT

Prentox

DDVP AEROSOL CONCENTRATE #G-1553

AN INSECTICIDE FOR FORMULATING USE

ACTIVE INGREDIENTS:

2,2-Dichlorovinyl dimethyl phosphate	33.33%
Related compounds	2.54%
1,1,1-Trichloroethane	64.13%
TOTAL:	100.00%

DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY PERFORATING OR CRUSHING. BURY OR DISCARD IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

NET CONTENT _____ GALLONS

DANGER KEEP OUT OF REACH OF CHILDREN

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN OR EYES. DO NOT GET IN EYES OR ON SKIN. DO NOT BREATHE FUMES. RAPIDLY ABSORBED THROUGH SKIN. IN CASE OF CONTACT, WASH IMMEDIATELY WITH SOAP AND WATER. DO NOT CONTAMINATE FEED AND FOODSTUFFS. WASH HANDS, ARMS AND FACE THOROUGHLY WITH SOAP AND WATER BEFORE EATING OR SMOKING. WASH ALL CONTAMINATED CLOTHING WITH SOAP AND HOT WATER BEFORE REUSE. WEAR NATURAL RUBBER GLOVES, WATER-PROOF PROTECTIVE CLOTHING AND GOGGLES. KEEP ALL UNPROTECTED PERSONS OUT OF OPERATIONAL AREA WHILE APPLYING THE INSECTICIDE, OR WORKING IN TREATED AREAS DURING POSTED PERIOD, WEAR A MASK OR RESPIRATOR OF A TYPE PASSED BY THE U.S. BUREAU OF MINES FOR PROTECTION AGAINST DIMETHYL DICHLOROVINYL PHOSPHATE.

Formulators using this product are responsible for providing data to support their Registration. Technical Bulletin available on request.

E.P.A. Reg. No. 655-317



SEE SIDE PANEL FOR ANTIDOTE & NOTE TO PHYSICIAN.

Prentiss Drug & Chemical Co., Inc.

PLANT 338 WILSON AVENUE, NEWARK, N. J. 07105

OFFICE 363 SEVENTH AVENUE, NEW YORK, N. Y. 10001

A PRENTISS PEST-TESTED PRODUCT

Prentox

ANTIDOTE:

If swallowed, give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep warm and quiet. Call a Physician immediately. In case of skin contact, remove contaminate clothing and immediately wash with soap and water.

NOTE TO PHYSICIAN:

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, Atropine Sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams until complete Atropinization has occurred. 2-PAM is also antidotal and may be administered in conjunction with Atropine.

Morphine is contraindicated.

Clear chest by postural drainage.

Observe patient continuously for 48 hours.

Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor.

Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.



Prentiss Drug & Chemical Co., Inc.

PLANT 338 WILSON AVENUE, NEWARK, N. J. 07105

OFFICE 363 SEVENTH AVENUE, NEW YORK, N. Y. 10001