

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not enter treated areas without protective clothing until dusts have settled. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: If container is damaged: Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with the pesticide and wear protective gear. Sweep the spill into a salvage drum. Dispose of wastes as below.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE (cont'd.)

Mix with pyrophyllite or talc. A pH of 6.5 to 7.0 is desirable. Not compatible with alkaline materials. Mix with carrier to give a Pyrethrin content of 0.1%, 0.15%, 0.2% and 0.3% (Equivalent to 5%, 7-1/2%, 10% or 15% of PYRONYL DRY CONCENTRATE)

The following dosage rates are recommended as guidelines, with variations on different types of crops, their age and locality.

Crop	Insects	Pyronyl Dry Content of Dust	Pounds Per Acre
Asparagus	Asparagus Beetle	5%	20-30 lbs
Beans	Aphids	5%	30 lbs
	Leaf Hopper (White Fly or Bean Jassid)	5%	20-30 lbs
	Mexican Bean Beetle	5-10%	25-35 lbs
	Thrips	5-10%	30 lbs
	Twelve Spotted Beetle	10%	45 lbs
Beets	Hawaiian Beet		
	Webworm	5-7.5%	20-30 lbs
	Webworm	5-10%	20 lbs
Broccoli	(See Cabbage)		
Brussels Sprouts	(See Cabbage)		
Cabbage	Cabbage Looper	7.5-10%	20 lbs
	Diamond Back Moth Larvae	7.5-10%	20 lbs
	Flea Beetle	10%	20 lbs
	Harlequin Bug	15%	30 lbs
	Imported Cabbage Worms	7.5-10%	20 lbs

prentox[®]
PYRONYL DRY
CONCENTRATE

1984

ACTIVE INGREDIENTS:

Pyrethrins	2.2%
Petroleum Distillates	37.8%
INERT INGREDIENTS:	60.0%
TOTAL:	100.0%

PRENTOX[®]—Registered Trademark of Prentiss Drug & Chemical Co., Inc.

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT

- If swallowed — Contact a physician immediately.
- If inhaled — Remove person to fresh air.
- If on skin — Remove contaminated clothing and wash affected areas with soap and water.
- If in eyes — Flush eyes with plenty of water.

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

If inhaled or swallowed: Keep out of reach of children. Avoid breathing dust. Wash thoroughly after handling prior to eating or smoking. Do not allow dust to contact food, feedstuffs or water supplies. Do not apply this product in a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

ENVIRONMENTAL HAZARDS


This product is toxic to fish. Do not apply directly to lakes, ponds, or streams. Apply this product as specified on the label.

E.P.A. REG. NO. 133-347

E.P.A. EST. NO.

NET CONTENTS

LOT #



Plant: Kaolin Road
 Sandersville, GA. 31082
 Office: C.B. 2000
 Clarksville, N.C. 27012

DIRECTIONS FOR USE

Crop	Insects
Cantaloupe	(See Melons)
Cauliflower	(See Cabbage)
Cherry	Flea Hopper Celery Leaf Tier Celery Looper
Collard	(See Cabbage)
Cranberries	Fire Worm Leaf Hopper
Cucumbers	Flea Beetle Spotted Cucumber Beetle Squash Bug Nymphs Striped Cucumber Beetle
Grapes	Leaf Hopper -- Nymphs Adults
Grass (Lawn)	Sod Webworm
Lettuce	(See Cabbage)
Melons	Squash Bug Nymphs Spotted Cucumber Beetle Striped Cucumber Beetle Thrips
Peppermint	Flea Beetle Western Twelve Spotted Cucumber Beetle
Potatoes	Aphids Colorado Potato Beetle Flea Beetle Leaf Hopper Psyllid
Radish	Flea Beetle
Spinach	Aphids Cabbage Looper Webworm
Squash	Mexican Bean Beetle Colorado Potato Beetle Squash Bug Nymphs Striped Cucumber Beetle
Tomatoes	Aphids Flea Beetle Psyllid Thrips
Watermelon	(See Melons)