

A PRENTISS PEST-TESTED PRODUCT

Prentox

PRENTISS MANUFACTURING COMPANY, PHILADELPHIA, PA.

655-423

ACTIVE INGREDIENT:	
2,2-Dichloropropionic acid, sodium salt of Dalapon ...	85%
(2,2-dichloropropionic acid equivalent 74%)	
INERT INGREDIENT:	<u>15%</u>
	TOTAL: 100%

CAUTION

KEEP OUT OF REACH OF CHILDREN

DO NOT USE IN NEARLY OPEN.
RETURN TO ORIGINAL CONDI-
TIONER, OR DESTROY BY
FRAGMENTING OR CRUSHING
AND BURYING IN A DEEP
HOLE.

AVOID CONTACT WITH EYES, SKIN OR CLOTHING.
USE ONLY AS DIRECTED. MAY CAUSE SKIN IRRI-
TATION. IN CASE OF CONTACT WITH EYES,
FLUSH WITH PLenty OF WATER AND GET MEDICAL
ATTENTION. IF ON SKIN, REMOVE CONTAMINATED
CLOTHING AND WASH SKIN WITH PLenty OF WATER.
WASH CLOTHING BEFORE REUSE.

Manufactured for:

Cameraman's Note
Poor Copy



... Reg. No. 655-423

Application:

This herbicide is used for the control of annual grasses, such as foxtail, crabgrass, and water grass, and perennial grasses, such as bermuda grass, Johnson grass, blue grass, quackgrass, and grass, cattails and phragmites.

Although this herbicide is absorbed through plant roots, it is not effective when applied as a foliar spray.

Best results are obtained when this herbicide is applied to vigorously growing grasses before the seed stalks have developed. Several weeks may elapse before maximum effect. If grass is not growing well, due to drought, cold or over maturity, results may be poor. The application should be repeated as necessary. Since its persistence is limited, new seedlings may develop in a treated area. Cultivate or retreat with a herbicide.

Precautions:

This herbicide will inhibit germination and plant growth for several months, depending on the application rate, soil type, soil moisture, rainfall, and temperature. It is advised that planting is delayed for 30 days after ploughing when using such crops as corn, beans, etc. Application is never in cold dry soil, and some time will be required before planting sensitive crops. Do not graze with dairy animals on vegetation treated with this herbicide.

Directions for Use:

Preparation:

Measure out spray tank mix at least 1/2 gallon, and add 1/2 gallon of water. Add sufficient water (10-40 gallons per acre), to give good coverage. Add 1/2 ounce of wetting agent, which is adequate for many uses. However, with certain perennial grasses that are difficult to wet, or when this herbicide is applied at low rates, especially in high dilution, 1/2 pint of non-ionic wetting agent per 1/2 gallon of spray, may be helpful in obtaining desired control.

... containing ... water ...

Application:

Use the suggested rate of application ... for the present treatment ...

Precautions:

When using spray boom, use 15 lbs. of herbicide per acre in at least 40 gallons of water. Repeat application in 2-4 weeks, and follow up with spot treatments. Use an orchard-type sprayer, use 10 lbs. of herbicide in 100 gallons of water, that foliage is wet thoroughly.

For the control of giant foxtail, no other grasses, in fence rows and other non-crop areas, use 1/2 gallon of herbicide in 20 gallons of water, spray to thoroughly wet the foliage, roots and rhizomes.

Precautions:

Do not use this herbicide in areas where it will be applied to sensitive crops. Do not use this herbicide in areas where it will be applied to sensitive crops. Do not use this herbicide in areas where it will be applied to sensitive crops.

Precautions:

Do not use this herbicide in areas where it will be applied to sensitive crops. Do not use this herbicide in areas where it will be applied to sensitive crops.

If treatment is not successful, repeat application ...

Manufacturer:
Dow Chemical Company