A PRENTISS PEST-TESTED PRODUCT

FOR USE IN FEDERALLY INSPECTED **MEAT & POULTRY ESTABLISHMENTS-**

DIRECTIONS FOR USE - DILUTE 1 PART PRENTOX PYRONYL OIL CONCENTRATE #3610 WITH 3 PARTS BASE OIL AND USE WITH A ULV SPRAYING DEVICE AT THE RATE OF ONE DUNCE PER THOUSAND CUBIC FEET OF ROOM SPACE. APPLY THE SPRAY INTO ALL CRACKS AND CREVICES IN WOOD-WORK, WALLS, FLOORS, UNDERNEATH SINKS, BEHIND PIPES AND IN ALL PLACES WHERE THESE INSECTS SHELTER. CON-TACT AS MANY INSECTS AS POSSIBLE. REPEAT DOSAGE AS NECESSARY FOR CONTROL OF ROACHES, SILVER-FISH, WATERBUGS, EARWIGS, ANTS, SPIDERS.

BEFORE USING, EXPOSED FOOD PRODUCTS AND PACKAGING MATERIALS MUST BE REMOVED FROM ROOM OR PROTECTED, AFTER USING, EQUIPMENT AND UTENSILS MUST BE THOROUGHLY RINSED WITH POTABLE WATER BEFORE OPERATIONS ARE RE-SUMED. USE ONLY WHEN PLANT IS NOT IN OPERA-

DO NOT POUR OR SPILL NEAR OPEN FLAME, MIX WELL IN CONTAINER BEFORE EACH USE. KEEP CON-TAINER CLOSED AND STORE IN A COOL PLACE.

DO NOT REUSE EMPTY DRUM, RETURN TO DRUM RECONDITIONER OR DESTROY BY PERFORATING OR CRUSHING, AND BURYING IN A SAFE PLACE.

APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

HARMFUL IF SWALLOWED. DO NOT APPLY DIRECTLY TO FOOD. COVER OR REMOVE ANY FOOD AND FOOD PROCESSING EQUIPMENT DURING APPLICATION, DO NOT APPLY WHILE FOOD PROCESSING IS UNDERWAY. AFTER SPRAYING IN MEAT PACKING PLANTS. BAKERIES AND OTHER FOOD PROCESSING PLANTS. WASH ALL EQUIPMENT, BENCHES, SHELVING, ETC. WHERE EXPOSED FOOD WILL BE HANDLED, WITH POTABLE WATER.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.



PYRONYL CONCENTRATE

ACTIVE INGREDIENTS:

TOTAL:	100.00%
Petroleum Distillates	81.00%
dicarboximide**	. 10.00%
N-octyl bicycloheptene	
Piperonyl Butoxide, Technical*	. 6.00%
Pyrethrins	

*Consists of 4.80% (Butylcarbityl) (6-propylpiperoonyl) Ether and 1,20% related compounds. **MGK-264 In secticide Synergist.

CAUTION KEEP OUT OF REACH CEMPTED

OF CHILDREN

NET CONTENTS _____

E. P. A. Est. No. 655-NJ-1

DIRECTIONS FOR USE

SPACE SPRAY - Use undiluted to kill accessible stages of insects* infesting dried fruit products in storage or being processed: Close room or warehouse and shut off all ventil-

ating systems. Use a ULV spraying device to disperse this

product as a fine mist in the air above the trays and shelves.

Do not apply to fruit directly. Use one ounce per 1000 cubic feet of space. Repeat application at this dosage in any given month. Leave the room closed for one hour. Do not re-

*Indian Meal Moth, Dried Fruit Beetle, Confused Flour Beetle, Saw-toothed Grain Beetle and Roaches.

SPACE SPRAY - Use undiluted to kill the adult and accessible forms of insect pests* of stored food (stored in multiwall paper bags or in cloth bags) in warehouses, stored

age rooms & similar locations: Close room or warehouse and

shut off all ventilating systems. Use a ULV spraying de-

vice to dispense this product. First direct spray into corners

vice to dispense this product. First direct spray into corners under pallets and around stacks of packaged goods hitting as many insects as possible and then treated as a space spray. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose. Use a dosage of one fluid ounce of this spray per 1000 cubic feet of room space. Repeat applications as necessary for insect control. Leaved the room closed for one hour. Do not remain in treated

*Rice Weevil. Yellow Meal Worm, Cadelle, Confused Flour

Beetle, Saw-toothed Grain Beetle, Angoumois, Grain Moth, Mediterranean Flour Moth and Indian Meal Moth.

SPACE SPRAY - Use diluted to kill flies, mosquitoes and

gnats in open areas: Dilute concentrate with Petroleum Distillate such as Klearol N.F. at the rate of 1 part concentrate to 2 parts oil and mix well. Use at a time when

the temperature is col (75. or less) and the wind velocity is approximately 5 MPH, with a ULV spraying device, spray

at a dosage rate of 15 ounces per minute up-wind of the area to be controlled. If the equipment is truck mounted,

keep the speed to approximately 5 MPH with a swath width of no more than 300 feet. Repeat application as required.

SPACE SPRAY AND CONTACT SPRAY - Use undiluted to

kill roaches, silverfish, waterbugs, earwigs, ants, spiders, in sheltered situations: With a ULV spraying device.

a dosage rate of approximately one-half fluid ounce per 1000 cubic feet or room space. Repeat dosage as neces-

sary for control. Leave the room closed for one hour. Do not remain in treated areas and ventilate area before re-

Do not allow spray to contact stored food. Use at

areas and ventilate area before re-entry.

main in treated areas and ventilate area before re-entry.

FUNGICIDE AND HODELLICIDE ACT. P.A. Reg. No. 655-450 FOR ECON. C



Prentiss Drug & Chemical Co., Inc.

PLANT: 338 WILSON AVENUE, NEWARK, N. J. 07105

OFFICE: 363 SEVENTH AVENUE, NEW YORK, N. Y. 10001