

BEST DOCUMENT AVAILABLE

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. If swallowed do not induce vomiting. Call a physician immediately.

Do not apply directly to food in commercial food handling facilities cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in meat packing plants, bakeries and other food processing plants wash all equipment, benches, shelving, etc. where exposed food will be handled with potable water. In the home all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets, birds and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not dump into prohibited areas.

PESTICIDE DISPOSAL - Pesticide spray mixture or use water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures.

CONTAINER DISPOSAL - Empty container or equivalent thereof must be triple rinsed and disposed of in accordance with applicable Federal, State or local procedures.

INDOOR USE AS A SPACE SPRAY UNPLUGGED - The use of this product in food processing or food handling establishments should be confined to the periods when the plants or facilities are not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Close windows and doors and shut off ventilating systems. Use a fogger or vaporizer adjusted to deliver an aerosol spray in droplets of 50 microns in diameter and 80% less than 30 microns. Apply at a rate of 1 ounce per 1,000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area behind partitions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary. Sweep up and destroy taken insects.

This method of application will control exposed, accessible stages of FLIES including FRUIT FLIES, MOSQUITOES, SMALL FLYING MOTHS, GNATS, WASPS, HORNETS, CLOVER MITES, CHEESE MITES, CHEESE SKIPPERS, BOXELDER BUGS, EARWIGS, ANTS and the following common stored product pests: GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, SAW TOOTHED GRAIN BEETLES, SPIDER BEETLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, ANGIOMOUS GRAIN MOTHS, MEDITERRANEAN FLOUR MOTHS, INDIAN MEAL MOTHS, TOBACCO MOTHS, YELLOW MEALWORMS, DARK MEALWORMS, GRAIN MITES, CADELLES and RED FLOUR BEETLES.

CONTACT AND SURFACE SPRAY - For the control of roaches, dilute 4 ounces to make one gallon of finished solution in oil. Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around bases of machinery, behind shelves and drawers. Spray roaches directly whenever possible. Repeat application as necessary.

DIRTY FRUIT PRODUCTS in storage or being processed for control of CONFUSED FLOUR BEETLES, DRIED FRUIT BEETLES, INDIAN MEAL MOTHS and SAW TOOTHED GRAIN BEETLES. Remove and destroy infested products. As noted above disperse this product as a fine mist in the air above the trays and shelves. Do not apply to fruit directly. Use one half ounce per 1,000 cubic feet of space. Repeat application at this dosage in a 7 day period. Leave the room closed for one hour. Do not remain in treated areas and ventilate the area before re-entry. Sweep up and destroy taken insects.

STORED FOOD - Stored in multi-wall bags or cloth bags in water-tight storage rooms, flour mills, grain elevators, granaries for control of ANGIOMOUS GRAIN MOTHS, CADELLES, CONFUSED FLOUR BEETLES, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS, RICE WEEVILS, SAW TOOTHED GRAIN BEETLES, YELLOW MEALWORMS, CHEESE MITES, CHEESE SKIPPERS, CIGARETTE BEETLES, GRANARY WEEVILS, GRANARY WEEVILS, DARK MEALWORMS, RED FLOUR BEETLES. Remove and destroy infested products. As noted above disperse this product first by direct spray into corners, under pallets and around stacks of packaged foods hitting as many insects as possible and then treat as a space spray.

Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose. Use a dosage of one half fluid ounce per 1,000 cubic feet of space. Do not remain in treated areas and ventilate area before re-entry. This application is accessible exposed stages of those stored product pests listed above. Sweep up and destroy taken insects.

INDOOR USE AS A SPACE SPRAY UNPLUGGED - For use in conventional mechanical fogging equipment to control FLIES, FRUIT FLIES, MOSQUITOES, SMALL FLYING MOTHS, AND GNATS. Cover or remove exposed food and food handling surfaces. Close room and shut off all air conditioning or ventilating system equipment. Dilute 1 part concentrate plus 29 parts of refined kerosene and mix well. Apply at the rate of 1 to 2 fluid ounces per 1,000 cubic feet of space. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

and ventilate area before re-entry. This application is accessible exposed stages of those stored product pests listed above. Sweep up and destroy taken insects.

**prentiss
Pyronyl Oil Concentrate #3610**

For Use by Trained Personnel Only

Authorized by U.S.D.A. for use in Federally inspected meat and poultry plants.

ACTIVE INGREDIENTS:

Pyrethrin	3.00%
Piperonyl Butoxide Technical*	6.00%
N-Octyl bicycloheptane dicarboximide	10.00%
Mineral Oil	69.00%
Petroleum Distillates	12.00%
TOTAL	100.00%

*Consists of 4.60% (Butylcarbityl) (6-Propylpiperonyl) Ether and 1.20% related compounds

Prentiss Pyronyl Oil Concentrate #3610 is effective in the control of the below-listed insects if the applicator follows the directions for use as set forth in this label.

ANGIOMOUS GRAIN MOTHS - ANTS - BOXELDER BUGS - CADELLES - CHEESE MITES - CHEESE SKIPPERS - CIGARETTE BEETLES - CLOVER MITES - CONFUSED FLOUR BEETLES - DARK MEALWORMS - DEER FLIES - DRIED FRUIT BEETLES - DRUGSTORE BEETLES - EARWIGS - FRUIT FLIES (DROSOPHILA) - FUNGUS GNATS - GNATS - GRAIN MITES - GRANARY WEEVILS - HORN FLIES - HORNETS - HORSE FLIES - HOUSE FLIES - INDIAN MEAL MOTHS - LICE - MEDITERRANEAN FLOUR MOTHS - MOSQUITOES - MUSHROOM FLIES - RED FLOUR BEETLES - RICE WEEVILS - ROACHES - SAW TOOTHED GRAIN BEETLES - SMALL FLYING MOTHS - SPIDER BEETLES - STABLE FLIES - TOBACCO MOTHS - WASPS - YELLOW MEALWORMS

KEEP OUT OF REACH OF CHILDREN

CAUTION

E.P.A. Reg. No. 655-450

E.P.A. Est. No. 655-GA-1

DIRECTIONS FOR USE (Cont'd)

INDOOR USE AS A SPACE SPRAY UNPLUGGED - For use in conventional mechanical fogging equipment to control FLIES, FRUIT FLIES, MOSQUITOES, SMALL FLYING MOTHS, AND GNATS. Cover or remove exposed food and food handling surfaces. Close room and shut off all air conditioning or ventilating system equipment. Dilute 1 part concentrate plus 29 parts of refined kerosene and mix well. Apply at the rate of 1 to 2 fluid ounces per 1,000 cubic feet of space. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

and ventilate area before re-entry. This application is accessible exposed stages of those stored product pests listed above. Sweep up and destroy taken insects.

INDOOR USE AS A SPACE SPRAY UNPLUGGED - For use in conventional mechanical fogging equipment to control FLIES, FRUIT FLIES, MOSQUITOES, SMALL FLYING MOTHS, AND GNATS. Cover or remove exposed food and food handling surfaces. Close room and shut off all air conditioning or ventilating system equipment. Dilute 1 part concentrate plus 29 parts of refined kerosene and mix well. Apply at the rate of 1 to 2 fluid ounces per 1,000 cubic feet of space. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

NET CONTENTS _____ gal

LOT # _____



Manufactured by
prentiss

Plant: Kaulin Road
Sandersville, GA 31082
Office: C.B. 2000

BOXELDER BUGS, CIGARETTE BEETLES, DROSOPHILA, etc. and mix well and apply as 100 of spray mixture to 750 sq. surfaces of bins and storage. Careful to treat cracks, crevices, handling and processing area equipment. Spray through equipment. Spray insects off. For insect protection, close before loading. Thoroughly cartons or containers before use. To control INDIAN MEAL MOTHS, apply to the surface of feet. Apply at monthly intervals continuing until early fall. Make before spraying and mix. As an aid in control of MOSQUITOES, use in resting places. ANIMAL QUARTERS USE: (Cont'd) 1000: As a space spray, dilute 1 part concentrate plus 29 parts of refined kerosene and mix well. Apply at the rate of 1 to 2 fluid ounces per 1,000 cubic feet of space. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

INDOOR USE AS A SPACE SPRAY UNPLUGGED - For use in conventional mechanical fogging equipment to control FLIES, FRUIT FLIES, MOSQUITOES, SMALL FLYING MOTHS, AND GNATS. Cover or remove exposed food and food handling surfaces. Close room and shut off all air conditioning or ventilating system equipment. Dilute 1 part concentrate plus 29 parts of refined kerosene and mix well. Apply at the rate of 1 to 2 fluid ounces per 1,000 cubic feet of space. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

INDOOR USE AS A SPACE SPRAY UNPLUGGED - For use in conventional mechanical fogging equipment to control FLIES, FRUIT FLIES, MOSQUITOES, SMALL FLYING MOTHS, AND GNATS. Cover or remove exposed food and food handling surfaces. Close room and shut off all air conditioning or ventilating system equipment. Dilute 1 part concentrate plus 29 parts of refined kerosene and mix well. Apply at the rate of 1 to 2 fluid ounces per 1,000 cubic feet of space. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

INDOOR USE AS A SPACE SPRAY UNPLUGGED - For use in conventional mechanical fogging equipment to control FLIES, FRUIT FLIES, MOSQUITOES, SMALL FLYING MOTHS, AND GNATS. Cover or remove exposed food and food handling surfaces. Close room and shut off all air conditioning or ventilating system equipment. Dilute 1 part concentrate plus 29 parts of refined kerosene and mix well. Apply at the rate of 1 to 2 fluid ounces per 1,000 cubic feet of space. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

- If swallowed** - Call a physician or Poison Control Center immediately. **DO NOT INDUCE VOMITING.** Vomiting may cause aspiration pneumonia.
- If inhaled** - Remove victim to fresh air. Apply artificial respiration if indicated.
- If on skin** - Remove contaminated clothing and wash affected areas with soap and water.
- If in eyes** - Flush eyes with plenty of water. Call a physician immediately.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

LEFT PANEL

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
Open dumping prohibited.

STORAGE: If container is damaged: STOP any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material, such as clay, diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. Dispose of on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Place any leaking container into a similar drum or glass container. Do not store, use pour or spill near heat or open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

PLASTIC CONTAINERS ONLY: DO NOT STORE IN CAR TRUNK WHERE TEMPERATURES MAY EXCEED 120°F.

300 / 104227
17 / 3

Prentiss Drug and Chemical
Co., Inc.
363 Seventh Avenue
New York, NY 10001

Gentlemen:

Subject: Pretox Pyronyl Oil Concentrate #3510
EPA Registration No. 655-450
Application Dated July 14, 1981

Your proposed amendment to bring the labeling for this product into compliance with PR Notice 83-3 is acceptable. A stamped copy of the draft labeling is enclosed for your records. Submit five (5) copies of the finished printed labeling before the product is released for shipment under the amended labeling.

Sincerely yours,

Timothy A. Gardner
Product Manager (17)
Insecticide-Acariocide Branch
Registration Division (TS-767)

Enclosure

TS-767: ~~TS-767~~ DCR-12110:WALG-1231C:lna:Raven:479-2013:11/9/83

CONCURRENTS							
SYMBOL	TS 767						
SURVEYED	Tanner						
DATE	11/11/83						