

prentox

Disinfectant Spray Concentrate #101

Authorized by U.S.D.A. for use in federally inspected meat and poultry plants

ACTIVE INGREDIENTS:

Pyrethrins	1.20%
Piperonyl Butoxide Technical	9.60%
Petroleum Distillates	81.20%
INERT INGREDIENTS	8.00%
TOTAL	100.00%

*Equivalent to 7.68% (Butylcarbityl) (6 propylpiperonyl) ether and 1.92% related compounds

ACCEPTED
with conditions
in 1933

11/27/1933

KEEP OUT OF REACH OF CHILDREN

CAUTION

655-452

655-452

N.Y.F.D.C. OF A. NO. 4146

CONTENTS gal. Lot #

FOR FLEAS, HOUSE FLIES, MOSQUITOES AND
OTHER PESTS AND DISINFECTANT FOR
MEAT AND POULTRY PLANTS

FOR DISINFECTING MEAT AND POULTRY
PLANTS AND OTHER BODIES WHERE
NECESSARY

FOR DISINFECTING MEAT AND POULTRY
PLANTS AND OTHER BODIES WHERE
NECESSARY

For effective control of BLATTELIDAE (FLIES), MUSCULIDAE (FLIES) AND (SUCTION)
BLOOD-SUCKING INSECTS (SHEEP TICKS AND LICE) on animals, apply a
part of the concentrate to the animal's body, paying special attention to
the head, neck, and between the legs.

To control POULTRY LICE, apply a part of the concentrate to the
body of the birds, paying special attention to the neck, head, and
between the legs.

For control of BEDBUGS AND BITES in poultry houses, apply a part of the
concentrate to the walls, floor, and other surfaces where the
bedbugs and ticks hide.

To control SHEEP TICKS AND LICE, apply a part of the concentrate to the
body of the sheep, paying special attention to the neck, head, and
between the legs.

Directions for Use (Continued)

To kill FLEAS ON LIVESTOCK AND PETS and to obtain protection against reinfestation
Dilute at a rate of 1 part concentrate and 9 parts water and bat the animal by dipping or
spraying. For best results against fleas and ticks on dogs, the harness or animal quarters
and bedding should also be treated.

To control SPIDERS AND TICKS Dilute at the rate of 1 part concentrate and 24 parts of
water and apply by individual treatment, spraying or pouring 1 to 2 ounces on each side of
large animals or a few drops into ears of dogs.

In horse milking parlors, wash rooms and dairies To control FLEAS, MOSQUITOES AND
GRATS Dilute at the rate of 1 part of concentrate and nine parts of water or oil and apply on
a rag or tow mat, directing spray toward the ceiling and upper corners until the area is filled
with mist using about 2 quarts per 1,000 cubic feet of space. For best results, close doors
and windows before spraying and keep them closed for one to fifteen minutes. Vacuum the
area and outside before re-cupping. Repeat application as necessary.

MOSQUITO CONTROL OUTDOORS To control ADULT MOSQUITOES using ground
application dilute 1 part concentrate and 7 parts water or oil. Apply on a tow mat or rag
Treat entire area with particular attention to shrubbery and vegetation where mosquitoes
may rest, and use of marshy areas, tree trunks and stagnant pools or ponds. Repeat or
necessary usually once or twice a week.

When mobile or truck-mounted equipment, the above application rate will result in a 400 to
500 ft. treated traveling at 15 mph per hour. Discharge spray at a rate of 40 gallons per
hour.

Application rate of 1 gallon of concentrate to 6 gallons of water
per acre.

To control MOSQUITO LARVAE Dilute at a rate of 1 part concentrate and 150 parts of
water in 50 gallons and apply at a rate of 20 to 25 gallons per acre. Apply a dose of water
to the water body and wash marshes by bodies of water where larvae may
breed. To control the dosage of 1000 may result.

ON GROWING CROPS May be used on outdoor and indoor greenhouses, etc. to
prevent the development of insects on a crop. Do not use harvest spray after
other chemicals have been used due to residue toxicity. When applied in or on a
crop, the concentrate may be used in combination with
other materials where resistance to a problem or insecticide is high. Do not
use on plants in contact with spray residue.

For the control of insects on plants, use 1 to 2 quarts per acre. The
water spray through coverage is about 3000 ft. Coverage of 1000 ft. covers
about 1/2 acre.

ASPARAGUS, BEANS, BEETROOTS, BELL PEPPERS, BRUSSELS SPROUTS, CABBAGE, CARROTS,
CAULIFLOWER, CELERY, CUCUMBERS, EGGPLANT, GREEN BEANS, GREEN PEAS,
KIDNEY BEANS, LIMA BEANS, MUNG BEANS, PEA, PUMPKIN, SPINACH,
SWEET CORN, TOMATOES, TURNIPS, ZUCCHINI.

BROCCOLI, CABBAGE, BRUSSELS SPROUTS, CAULIFLOWER, CARROTS, CUCUMBERS,
EGGPLANT, GREEN BEANS, GREEN PEAS, KIDNEY BEANS, LIMA BEANS,
MUNG BEANS, PEA, PUMPKIN, SPINACH, SWEET CORN, TOMATOES,
TURNIPS, ZUCCHINI.

CELERY, CUCUMBERS, EGGPLANT, GREEN BEANS, GREEN PEAS,
KIDNEY BEANS, LIMA BEANS, MUNG BEANS, PEA, PUMPKIN,
SPINACH, SWEET CORN, TOMATOES, TURNIPS, ZUCCHINI.

LETTUCE, PEAS, PUMPKIN, SPINACH, SWEET CORN, TOMATOES,
TURNIPS, ZUCCHINI.

MUSHRUM GREENS, RICE, COLLARDS, TURNIPS, ASPARAGUS, BEANS,
BROCCOLI, CABBAGE, BRUSSELS SPROUTS, CAULIFLOWER,
CARROTS, CUCUMBERS, EGGPLANT, GREEN BEANS, GREEN PEAS,
KIDNEY BEANS, LIMA BEANS, MUNG BEANS, PEA, PUMPKIN,
SPINACH, SWEET CORN, TOMATOES, TURNIPS, ZUCCHINI.

SPINACH, CABBAGE, CARROTS, CUCUMBERS, EGGPLANT,
GREEN BEANS, GREEN PEAS, KIDNEY BEANS, LIMA BEANS,
MUNG BEANS, PEA, PUMPKIN, SPINACH, SWEET CORN,
TOMATOES, TURNIPS, ZUCCHINI.

TOMATOES, TURNIPS, ZUCCHINI.

TURNIPS, ZUCCHINI.

ZUCCHINI.

1. Apply concentrate to plants and fruit in baskets on the ends and plants. Use spray of
right pressure for applying at the rate of 1 to 2 quarts of the diluted spray to a bush to
lead to complete. Direct the spray for a maximum coverage of the baskets or plants.
It is important to spray between and beneath the branches. The spray will spray
the leaves, but the emulsion remains on the dead parts so that they are ready to be
removed.

2. Spray the row ends stacked in the field.

3. Use baskets in the packed cases after dumping the fruit and after it is well
packed.

4. After washing and cleaning up the inside of the processing plant and after
draining fruitless, the entire floor inside the cannery should be sprayed at a dilution
of 1 part concentrate to 50 parts water (1 part to 1 gallon) 1 part water to 1
gallon concentrate with a water at the rate of 1 gallon to 750 square feet direct on by a
40 ft. nozzle and then, strong special attention to forcing spray into all cracks and
crevices.

DOCUMENT AVAILABLE

prentiss Plant: Kaolin Road
Sandersville, GA 31082
Office: 263 7th Avenue

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

- If swallowed - Call a physician or Poison Control Center immediately. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia.
- If inhaled - Remove victim to fresh air. Apply artificial respiration if indicated.
- If on skin - Remove contaminated clothing and wash affected areas with soap and water.
- If in eyes - Flush eyes with plenty of water. Call a physician immediately.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

BEST DOCUMENT AVAILABLE

LEFT PANEL

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
Open dumping prohibited.

STORAGE: If container is damaged: STOP any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required as prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material, such as clay, diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. Dispose of on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Place any leaking container into a similar drum or glass container. Do not store, use pour or spill near heat or open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

CONTAINER DISPOSAL. Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

PLASTIC CONTAINERS ONLY: DO NOT STORE IN CAR TRUNK WHERE TEMPERATURES MAY EXCEED 120°F.

Frontiss Drug & Chemical Co. Inc.
363 7th Avenue
New York, NY 10001

Gentlemen:

Subject: Prentox Emulsifiable Spray
EPA Registration No. 655-452
Your Submission Dated July 15, 1983

655 452
300 / 104555
17 3

The amendment referred to above, submitted in connection with registration under FIFRA section 3(c) (7) (B), is acceptable, provided that you:

1. Make the following changes before you release the product for shipment bearing the amended labeling:
 - a. Delete the words "Keep out of lakes, streams, and ponds." Replace with "Do not add directly to water."
 - b. Delete the words "Apply artificial respiration if indicated."
 - c. Add the words "if irritation persists" after the words "Call a physician immediately" in the "If, in eyes" statement of practical treatment.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

Timothy A. Gardner
Product Manager (17)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

TS-767:GARDNER:DCP-31547:WANG-0385C:D.D:Raven:479-2013:9/15/83

(CAS)

CONCURRENTS							
SYMBOL							
SUBNAME							
DATE							