prentox

VAPBAY CONCENTRATE

An insecticide concentrate for formulating use only.

ACTIVE INGREDIENTS:	
0-Isopropoxyphenyl Methylcarbamate*	20.0%
2, 2-Dichlorovinyl Dimethyl Phosphate	10.0%
Related Compounds	0.8%
INERT INGREDIENTS:	69.2%
TOTAL:	100.0%

*

KEEP OUT OF REACH OF CHILDREN

DANGER
POISON

DIRECTIONS FOR USE

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

POISONOUS IF SWALLOWED INHALED OR ABSORBED THROUGH SKIN OR EYES. DON'T GET ON SKIN OR IN EYES. DO NOT BREATHE MIST. WEAR NATURAL RUBBER GLOVES, WATER PROOF PROTECTIVE CLOTHING AND GOGGLES WHEN HANDLING. WEAR A RESPIRATOR IDENTIFIED APPROVED BY THE MINING ENFORCEMENT AND SAFETY ADMINISTRATION (FORMERLY THE U.S. BUREAU OF MINES) AND BY THE NATIONAL INSTITUTE FOR OCCUPATIONAL SAFETY AND HEALTH UNDER THE PROVISIONS OF 30 CFR PART 11 FOR PROTECTION AGAINST THIS
IN CASE OF CONTACT WASH IMMEDIATELY WITH SOAP AND WATER. WASH HANDS, ARMS AND FACE THOROUGHLY WITH SOAP AND WATER BEFORE EATING OR SMOKING. WASH ALL CONTAMINATED CLOTHING WITH SOAP AND WATER BEFORE REUSE.
NOT FOR USE OR STORAGE IN OR AROUND THE HOME.
THIS PRODUCT IS TOXIC TO FISH, BIRDS AND OTHER WILDLIFE. DO NOT CONTAMINATE LAKES, STREAMS OR ESTUARIES. DISPOSAL IN ACCORDANCE WITH A NATIONAL POLLUTION DISCHARGE ELIMINATION SYSTEM PERMIT MAY BE UNLAWFUL. CONTACT YOUR REGIONAL OFFICE OF THE ENVIRONMENTAL PROTECTION AGENCY.

STATEMENT OF PRACTICAL TREATMENT

If swallowed — Call a physician immediately. Gastric lavage is not advised. If material was taken internally DO NOT INDUCE VOMITING. Most effective treatment is not available, limiting [illegible] to induce vomiting. Give large quantities of water and inert absorbent [illegible]. Repeat until vomiting ceases. Do not give anything by mouth to unconscious person. **If inhaled** — [illegible] Apply respiration if indicated. **If on skin** — Wash contaminated skin with soap or detergent and water. **If in eyes** — Flush eyes with plenty of water for at least 15 minutes. Get medical attention promptly.

ANTIDOTE & FIRST AID TREATMENT

ATROPINE [illegible] ANTIMETABOLYN ACETANILIDE [illegible]
IF ANY NERVOUSNESS, MUSCULAR TWITCHING, Atropine [illegible] SPECIFIC ANTIDOTE UNLESS WARNING SYMPTOMS APPEAR. See WARNING SYMPTOMS below.
Keep patient [illegible] and [illegible] artificial respiration if [illegible]. Transport the patient immediately to the nearest [illegible].

NOTE TO PHYSICIAN —
SYMPTOMS Symptoms include weakness, headache, tightness of chest, blurred vision, [illegible] pupils, excessive salivation, sweating, nausea, vomiting, diarrhea, abdominal cramps.
TREATMENT Atropine is the specific therapeutic antidote against parasympathetic stimulation. Atropine Sulfate should be injected at 5-minute intervals in doses of 1/50 to 1/20 grain until complete Atropinization has occurred. 2-PAM is also an antidote. Initial dose of 1 gram. Atropine [illegible] Morphine is contraindicated. Check test by postural drainage [illegible] several times a day.
Repeated exposure to cholinesterase inhibitors may without warning cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

STORAGE AND DISPOSAL:

PESTICIDE DISPOSAL [illegible]
CONTAINER DISPOSAL 30 & 55 gallon drums [illegible]
SMALLER CONTAINERS [illegible]
GENERAL [illegible]

[illegible] is a registered trademark of Farbenfabriken Bayer A.G. Leverkusen

EPA REG 655-454 **EPA EST NO 655-NJ1**

NET CONTENTS

LOT #

ACCEPTED
JUL 07 1981
For the Federal [illegible]

Manufactured by
prentiss
Drug & Chemical Co., Inc.
Plant: 338 Wilson Avenue
Newark, N.J. 07105
Office: 363 [illegible] Avenue
New York, N.Y. 10001