

prentox. DIAZINON 50W Insecticide

FOR AGRICULTURAL/
COMMERCIAL USE ONLY

E.P.A. Reg. No. 888-488
E.P.A. EST. NO. 894-NJ-2

Read these entire directions. Use strictly in accordance with label warnings and directions.

For control of certain insects on fruits, nuts, vegetables,
leafy crops, and ornamentals.

ACTIVE INGREDIENTS

DIAZINON (20-chloro-2-(diisopropyl 6-methyl-4-pyridinyl) phosphorothioate) 80%

INERT INGREDIENTS

..... 20%

TOTAL

..... 100%

DIAZINON 50W IS A WETTABLE POWDER.

NOTE TO USER — Some uses require more than 1 x 4 lb. bag. Make sure to read directions before purchasing. Please see attachment for directions.

PRENTOX® — Registered Trademark of Prentiss Drug & Chemical Co., Inc.

**KEEP OUT OF REACH OF CHILDREN
WARNING — AVISO**

"If used on the label, no use extra products have due to adequate label and explicit instructions" *655-456*

STATEMENT OF PRACTICAL TREATMENT

If Inhaled — Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching back of the throat with finger. Repeat until vomit is clear. Do not induce vomiting or give anything by mouth to an unconscious person.
If Irritated — Remove victim to fresh air and apply artificial respiration if indicated.
If In Eyes — Wash promptly with soap and water. Rinse thoroughly.
If In Ears — Rinse eyes with plenty of water and call a physician immediately.
NOTE TO PHYSICIAN — If signs of cholinesterase inhibition are present. Atropine or antidotal 2-PAM may be administered in conjunction with a atropine.

PRECAUTIONARY STATEMENTS

Hazardous to Humans and Domestic Animals

May be fatal if swallowed. May be deadly if inhaled. Do not breathe dust or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating or smoking. Keep out of reach of children and pets. Do not use on humans or household pets. Do not permit children or pets to play in treated areas until spray has completely dried.

Environmental Hazards
This product is toxic to fish, birds and/or other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, swamps, marshes and estuaries (except for recreational purposes and human and livestock consumption). Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish and other aquatic life are important resources. This product must not be used in areas where impact on threatened and endangered species is likely. Contact your State Fish and Game Agency for information pertaining to areas having threatened and endangered species. Apply this product only as specified on this label.

This product is highly toxic to bees used to direct treatment or to residues remaining in the treated area. Do not apply when bees are active in the crop cover or on weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

IMPORTANT:

Read the entire directions for use before using this product.

STORAGE & DISPOSAL

Do not contaminate water, food, feed, storage or disposal. Do not dump or pour into water.
STORAGE — If container is damaged, do not use. Do not use if contents are not in original container. Do not use if contents are not in original container. Do not use if contents are not in original container.
PESTICIDE DISPOSAL — Wastes resulting from use of this product may be disposed of by incineration at an approved waste disposal facility.
CONTAINER DISPOSAL — Completely empty bag. Do not reuse. Do not reuse. Do not reuse.

GENERAL INFORMATION

This product is a systemic insecticide. It is effective against a wide range of insects. It is effective against a wide range of insects. It is effective against a wide range of insects.

FRUIT AND NUT INSECTS

Almonds — Mites, Two-spotted Spider Mites, Leafhoppers, Peach Tree Weevil, etc.
Apples — Leafhoppers, Aphids, Pear Leaf Miner, etc.
Berries — Leafhoppers, Aphids, etc.
Cherries — Leafhoppers, Aphids, etc.
Citrus — Citrus Red Spider Mite, etc.
Corn — European Corn Borer, etc.
Cucumbers — Aphids, etc.
Grapes — Grape Leaf Miner, etc.
Leafy Crops — Leafhoppers, Aphids, etc.
Peaches — Peach Tree Weevil, etc.
Pears — Pear Leaf Miner, etc.
Peanut — Peanut Aphid, etc.
Plum — Plum Aphid, etc.
Potatoes — Colorado Potato Beetle, etc.
Strawberries — Strawberry Leaf Miner, etc.
Tomatoes — Tomato Pinworm, etc.
Walnuts — Walnut Caterpillar, etc.

Beans — Pacific Spider Mite, Leafhoppers, Grape Berry Mite, Grape Leaf Miner, etc.
Berries — Leafhoppers, Aphids, etc.
Cherries — Leafhoppers, Aphids, etc.
Citrus — Citrus Red Spider Mite, etc.
Corn — European Corn Borer, etc.
Cucumbers — Aphids, etc.
Grapes — Grape Leaf Miner, etc.
Leafy Crops — Leafhoppers, Aphids, etc.
Peaches — Peach Tree Weevil, etc.
Pears — Pear Leaf Miner, etc.
Peanut — Peanut Aphid, etc.
Plum — Plum Aphid, etc.
Potatoes — Colorado Potato Beetle, etc.
Strawberries — Strawberry Leaf Miner, etc.
Tomatoes — Tomato Pinworm, etc.
Walnuts — Walnut Caterpillar, etc.

BEST DOCUMENT

655-456

655-456

BEST DOCUMENT AVAILABLE

FRUIT AND BERRY INSECTS

Almonds - Mites, Tort Borer. Apply 1/2 lb. per 100 gals. of water.
Panama Scales, San Jose Scales, Mite 1 lb. of Diazinon 50W plus 2 3/4 gals. dormant oil per 100 gals. of water and apply as a dormant spray.

Apples - Leafhoppers. Apply 1/2 lb. per 100 gallons of water in 3-5 applications beginning at petal fall and continuing through the growing season or apply as insecticidal beverage.
 San Jose Scales, Mealybugs, Apple Aphid eggs, European Red Mite eggs, Brown Mite eggs, Mite 1 lb. of Diazinon 50W plus 1 gal. of superior type oil or 2 gals. of dormant oil per 100 gals. of water. Apply as a dormant or a delayed dormant spray.
 Spotted Bud Moths, Fruit Tree Leaf Rollers, Mealybugs, Mite 1 lb. per 100 gals. of water. Make 1/2 lb. dormant spray applications beginning in the pink stage.
 Do not apply within 14 days of harvest.

Apples and Pears - Codling Moths, Rosy Apple Aphids, Green Apple Aphids, Woolly Apple Aphids, Mites. Mix 1 lb. of Diazinon 50W per 100 gallons of water. Make 3 to 8 cover spray applications beginning with the first cover. Applications should be timed according to insect infestation.

Apple Maggots - Mix 1 lb. of Diazinon 50W per 100 gallons of water. Make at least 3 cover spray applications when flies are active and laying eggs.

San Jose Scale Crawlers, Forster Scale Crawlers - Mix 1 lb. per 100 gallons of water. Apply early in the season when crawlers first appear. If infestation warrants make a second application 10-14 days later.

Nitrogen - Mite. Mix 1 lb. of Diazinon 50W in the regular cover spray. Do not spray on varieties of apples such as Golden Delicious which are susceptible to apple scab. Do not apply to apples or pears within 14 days of harvest.

Apricots - San Jose Scales, Brown Apple Scales, Walnut Scales, Apricot Mealybug, Mite. Mix 1 lb. of Diazinon 50W plus 2 gallons of dormant oil per 100 gallons of water. Apply as a dormant spray.

Aphids, Brown Mites (Clover Mites), Two-Spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers - Mix 1 lb. per 100 gallons of water. Apply when aphids, mites and scale crawlers first appear.

Olive Scale Crawlers - Mix 1 lb. of Diazinon 50W plus 1/2 gallon of light medium horticultural oil per 100 gallons of water. Apply when scale crawlers are present.

Apricot Mealybugs - Mix 1 lb. per 100 gals. of water. Apply as a cover spray from petal fall to June.

Do not apply within 10 days of harvest.

Raspberries - Cranberry Fruitworms, Cherry Fruitworms, Blueberry Maggots, Aphids, Thrips, Two-spotted Mites. Mix 1 lb. per 100 gals. of water. Apply up to 200 gals. of spray per acre or apply 2 lbs. per acre in sufficient water for thorough coverage.
 Do not apply within 7 days of harvest.

Cranberries - L. Blueberry, Raspberry, Blueberry, Leafhoppers, Raspberry - Leafhoppers, Thrips, Aphids, Two-spotted Mites, Raspberry Sawflies. Mix 1 lb. per 100 gals. of water and apply up to 200 gals. of spray per acre, or apply 2 lbs. per acre in sufficient water for thorough coverage.

Raspberry Fruitworms - Apply at the above rates when blossom buds separate and repeat application immediately before blossoms open.

Dryberry Mites - Mix 2 lbs. per 100 gals. of water. Apply as a dormant spray in early spring when canes are still on the ground and buds are closed. Make a second application when the buds are well developed but still closed.

Raspberry Crown Borer (Raspberry Root Borer) - Mix 4 lbs. per 100 to 200 gals. of water and apply as a drench to the crown area and to the lower canes. Apply either in the spring before borers begin to form, or at the fall after harvest.

Do not apply within 7 days of harvest.

Cherries - Brown Apple Scales, San Jose Scales, Black Scales, Black Cherry Aphid eggs, Brown Mite eggs. Mix 1 1/2 lbs. of Diazinon 50W plus 3 gals. of dormant oil per 100 gals. of water. Apply as a dormant spray.

Eye-spotted Bud Moths, Fruit Tree Leaf Rollers - Mix 1 lb. per 100 gals. of water. Apply in cover spray as necessary beginning with petal fall.

Leafhoppers - Apply 1 lb. per 100 gallons of water in 3-5 applications beginning at petal fall and continuing through the growing season or apply as insecticidal beverage.

Cherry Fruit Flies - Mix 1/2-1 lb. per 100 gals. of water. Make 3 or 4 applications beginning when adult flies begin to emerge and repeating at 10-day intervals.

Black Cherry Aphids - Mix 1 lb. per 100 gals. of water. Apply as a cover spray when aphids appear and repeat after 10 days.

San Jose Scale Crawlers - Mix 1 lb. per 100 gals. of water. Apply during the pre-pink stage or when crawlers first appear.

Cherry Root Mites - Mix 1 lb. per 100 gals. of water. Apply as a supplemental spray after harvest.

Do not apply within 10 days of harvest.

Citrus - Black-headed Fireworms, Rhinoceros Beetle. Apply 4 lbs. in 400 gallons of water per acre.

Cranberry Fruitworms - Apply 6 lbs. in 400 gallons of water per acre.
 Do not apply within 5 days of harvest.

Figs - Vinegar Flies (Drosophila spp.), Dried Fruit Beetles. Apply 1 lb. per 100 gals. of water.
 Do not apply within 5 days of harvest.

Figs - Filbert Leaf Rollers, Aphids. Apply 2-4 lbs. in 250-400 gals. of water per acre as a thorough cover spray.

Citrus Fruits - (Grapefruit, Lemons, Oranges, Tangerines, Tangelos, Citrus, Citrus, Lemons, and hybrids of same) - Citrus Aphids, Cottony-cushion Scale Crawlers and Soft Scale Crawlers. Apply 1 lb. per 100 gals. of water (maximum 1000 gals. per acre) as a thorough cover spray when aphids and crawler stages of scale insects first appear.

Fruit Tree Leaf Rollers - Mix 1/2-1 lb. per 100 gals. of spray (maximum 1000 gals. per acre when an average of 50 percent of eggs in egg masses have hatched in the spring).

Citrus Thrips - Apply 1 lb. in 100 gals. of water (maximum 1000 gals. per acre) when thrips first appear.

Citrus Snow Scales - Apply 1-2 lbs. per 100 gals. water (maximum 1000 gals. per acre) as a post-bloom spray and again as a summer spray. For heavy snow scale infestations use the higher dosage rate. Thorough coverage of tree trunk, limbs and twigs is essential. Do not apply during bloom.

Do not apply to citrus within 21 days of harvest.

Potatoes - Aphids, Flea Beetles, Colorado Potato Beetle, Colorado Potato Flea Beetle, Colorado Potato Flea Beetle, Colorado Potato Flea Beetle. Apply 1 lb. per acre.
 Do not apply within 30 days of harvest.

Peas - Pea Aphids, Flea Beetles, Colorado Potato Beetle, Colorado Potato Flea Beetle. Apply 1 lb. per acre.
 Do not apply within 10 days of harvest.

Spinach and Beets - Aphids, Dotted Leaf Miner. Apply 1 lb. per acre.
 Do not apply to spinach within 10 days or to beets within 14 days of harvest.

Squash (Summer and Winter) - See Melons and Squash above.

Bean Cans - Corn Earworms. Apply 2 1/2 lbs. per acre when first application is made. Repeat applications at 2-3 day intervals will usually be necessary. Beans to be picked immediately following last application. Cornborer may be detected by and on cattle and sheep.

Refer also to Corn Under Field and Forage Insects.

Tomatoes - Aphids, Dotted Leaf Miner. Apply 1 lb. per acre. In control of miners weekly applications are usually required.

Barred Cucumber Beetles, Fall Armyworms, Southern Armyworm - Apply 1 lb. per acre.

Vinegar Flies (Drosophila spp.) - Apply 1 lb. per acre. Weekly applications are usually required.

Do not apply within 1 day of harvest.

Watermelon (Honey) Beetles - Cyclamen Mites. Apply 1 lb. in 100 gals. of water per acre. Do not apply to watermelon during top of plants. Infestation is heavy. Make 4 or 5 applications at 10-14 day intervals after the first application. Repeat application as necessary.

Do not apply within 5 days of harvest.

655-456

VEGETABLE SOIL INSECTS

Diazinon 50W should always be applied as a spray in sufficient water to assure thorough coverage of the soil.

Carrots - Apply 4 lbs. per acre as a drench to the seed furrow at planting time.

Cauliflower (Florets and Subterranean) - Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Cauliflower, Celery, Celeriac, Cucumbers, Endive (escarole), Kale, Lettuce, Lima Beans, Mung Beans, Parsley, Peas, Peppers, Potatoes, Snap Beans, Spinach, Summer Squash, Sweet Corn, Sweet Chard, Tomatoes, Watermelons, Winter Squash - Broadcast 4-8 lbs. per acre just prior to planting. Work into the soil immediately 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.

Corn (Sweetcorn) - Broadcast - Cabbage, Carrots, Corn, Lettuce, Peas, Pole Beans, Potatoes, Snap Beans, Tomatoes, Turnips - Broadcast 2 lbs. per acre just prior to planting. Work into soil 4-8 inches immediately. This pre-plant treatment will also control other soil insects such as cutworms (surface and subterranean), mole crickets, root maggots and rootworms.

Other Crops - Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Celeriac, Endive (escarole), Kale, Lettuce, Mung Beans, Peppers, Potatoes, Tomatoes, Turnips - Broadcast 2 lbs. per acre when insects are present. Make application to plant base or holes 1-2 days prior to sowing of seed or transplanting, and work into soil 1-2 inches immediately.

Bean Maggots - Onions, Parsley. Application - Apply 2 lbs. per acre in sufficient water to drench the seed furrow at planting time.

Broadcast Application - Broadcast 4-8 lbs. per acre just prior to planting. Work into soil 3-4 inches immediately.

Root Maggots - Broccoli, Brussels Sprouts, Cabbage and Cauliflower. Broadcast Application - Broadcast 4-8 lbs. per acre and immediately work into the soil 3-4 inches before sowing, either in the seedbed or in the field or hole's transplanting.

Transplant Water (Drench) Application - Mix 1/2 lb. in 50 gals. of transplant water when 200-300 gals. of water are used per acre. May be applied at the rate of 1/2 cupful per plant either by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.

Rootworms - Mix 2 lbs. in 50 gals. of water. Apply 50-100 gals. per acre (1-1 1/3 gals. per 1,000 linear feet of row) in the seed furrow at planting time.

Watermelons - Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Carrots, Cauliflower, Celery, Celeriac, Cucumbers, Endive (escarole), Kale, Lettuce, Lima Beans, Mung Beans, Parsley, Peas, Peppers, Snap Beans, Spinach, Summer Squash, Sweet Corn, Sweet Potatoes, Sweet Chard, Tomatoes, Watermelons, Winter Squash - Broadcast 4-8 lbs. per acre just prior to planting, and work into soil 4-8 inches immediately.

Potatoes - Broadcast 6-12 lbs. per acre just prior to planting. Immediately disk or harrow into soil 4-8 inches.

Southern Field Fleas - Potatoes (Southeastern U.S.) For potatoes on mineral soils broadcast 4-8 lbs. per acre just prior to planting and immediately work into soil 3-6 inches depending on location of wireworms.

SOIL AND FORAGE INSECTS

Diazinon 50W should always be applied as a spray in sufficient water to assure thorough coverage of the foliage or soil depending on the type of application.

Alfalfa and Clover - Spotted Alfalfa Aphids, Yellow Clover Aphids, Pea Aphids, Leafhoppers, Plant Bug, Nymphs, Mites. Apply 1/2-1 lb. per acre.

Grasshoppers - Apply 1 lb. per acre.

Cutworms (Surface and Subterranean) - Broadcast 4-8 lbs. per acre just prior to planting. Work into the soil immediately 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.

Dairy and beef cattle and sheep (1) may be grazed immediately following application. (2) may be fed green alfalfa or clover immediately following cutting, and (3) may be fed hay provided alfalfa or clover is not cut for hay for 7 days following treatment.

Alfalfa Weevil Larvae - For control in the spring, apply 2-3 lbs. per acre when feeding damage to growing tops becomes evident. Repeat application if necessary. If damage is already extensive, cut the hay and make application to the stubble. Do not graze dairy and beef animals or sheep for 2 days. Do not cut for hay for 10 days following treatment.

Do not spray livestock directly, except as directed below under Livestock Insects with either air or ground equipment.

Do not treat during bloom to avoid injury to pollinating insects.

Corn - Corn Rootworm Larvae. Post-emergence basal treatment: Apply 1.4 to 2.0 oz. per 1,000 linear feet of corn row in sufficient water to wet base of corn plant and narrow band of soil next to the plant. Refer to row spacing chart for amounts recommended per acre. Give treated band with 2-3 inches of soil using a disk harrow or cultivator immediately after spraying. Application should be made at the first sign of corn rootworm larvae. Application should not be made later than June 24 in the Middle Corn Belt or July in the Northern Corn Belt.

ACCEPTED

FEB 10 1984

Under the Federal Insecticide, Fungicide, and Rodenticide Act, or amended, for the pesticide registered under EPA Reg. No. 655-456

Spider Mites, Clover Mites, Spider Mites, Dip - Root Weevils, European Pine Shoot Moths, European Red Mites, Fruit Beetles, Holly Blue Moths, Leafhoppers, Oblique-banded Leafhopper, Pear Leaf Miner, Prickly Pear Cactus, Cottony Cuckoo, Lecanium, Pine Needle, San Jose Soft Scale, Thrips, Two-spotted Mites, Webworms, Whiteflies Apple and Thorn Skeletonizers, Coloneaster Web- worms, Fall Webworms, Hemlock Cherries, Oak Curculionids, Oblique-banded Leafhoppers, Pear Slugs, Pear Caterpillars	0.7	1.5
Apple and Thorn Skeletonizers, Coloneaster Web- worms, Fall Webworms, Hemlock Cherries, Oak Curculionids, Oblique-banded Leafhoppers, Pear Slugs, Pear Caterpillars	1-32	3.25

Do not use on certain Ferns, Poinsettia, Hibiscus, Papaya, Pine, and Gardenia, since injury to the plants may occur.
 Mimosa Webworms: Mix 1/2 oz. in 3 gals. of water (or 1 lb. in 100 gals. of water) and apply as a thorough foliar spray when adults first appear. Make a second and third application at 4-5 week intervals.

LAWN INSECTS

To control insects listed below, apply recommended rates. For insects such as Ants, Brown Dog Ticks, Clover Mites, Crickets, Earwigs, Fleas, Millipedes, Scabbards, and Centipedes (Springtails) apply 1/2 oz. of Diazinon 50W in 100 gals. of water. For house flies, apply 1/2 oz. of Diazinon 50W in 100 gals. of water around the house as well as the house foundation wall to a height of 2-3 ft. Repeat application if necessary.

Insects	Gals./1,000 sq. ft.		Remarks
	Minimum	Maximum	
Lawn Chinch Bugs	2-3 oz. or 3-6 oz.	25" gals.	Use higher rate for longer residual control and in areas of dense growth such as St. Augustine lawns.
Ants, Armyworms, Clover Mites, Collembola (Springtails), Crickets, Cutworms, Digger Wasps, Earwigs, Fireflies, Lawn Billbugs, Sod Webworms (Lawn Moths), Softbugs	4 oz.	3 gals.	For lawn billbugs, irrigate the lawn thoroughly immediately after treatment. For fireflies, mow grass and water lawn before treatment but delay application if the grass is dry. Do not water again until necessary.
Brown Dog Ticks, Bermuda Mites, Chiggers, Fleas, Leafhoppers	1-2 oz.	3 gals.	Do not apply to animals. For brown dog ticks, spray grass and under shrubbery particularly near house.
Millipedes	8 oz.	3 gals.	
Rhodes-grass Scales	5 oz.	25" gals.	Apply when crawlers first emerge.

*Alternatively, make applications in 3 gals. of water per 1,000 sq. ft. and thoroughly water to facilitate following application.

Beetle Bugs: Spray bugs outside the home with 1/4-2/3 oz. in 1 gal. of water. At the same time, spray thoroughly around foundation, windows, doors, tree trunks, and other cracks and other places where the insects hide.

SEED TREATMENT — PLANTING BOX TREATMENT

Corn, Seedcorn beetles: Mix 3 oz. per bushel of seed as a dry mixture in the planter box as a seed treatment prior to planting.

Corn, Plant, Soybeans, and Beans (Lima, Bush, White, and Kidney)

Seedcorn mites: Mix 1/2 oz. per bushel of seed as a dry mixture in the planter box as a seed treatment prior to planting.

Planter-box mixing instructions: Stir seed steadily using a mixing stick. Additional mixing will result from the operation of the planter. **DO NOT MIX WITH BARE HANDS.**

Note: If seed has not been treated with a fungicide, use Captan or Thiram in combination with Diazinon 50W at rates indicated on fungicide package. Failure to use a fungicide with Diazinon 50W may result in injury to the seed. If the seed treatment increases the friction between seeds sufficiently to reduce the rate of delivery through the planter, 1/2 oz. of graphite per bushel of seed should be added as a draft lubricant. Adjust the planter to give the required rate of seed delivery with the treated seed.

Warning: Treated seed must not be used for or mixed with food or animal feed, or processed for it. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time. Exposed treated seeds may be hazardous to birds and other wildlife. Dispose of all excess treated seeds by burial away from bodies of water.

FLY CONTROL

Fly control in barns and animal sleeping quarters (except dairy barns, milking parlors, and poultry houses):

Residual Sprays: Remove animals from buildings prior to treatment and keep animals out for at least four hours. Mix 2 lbs. of Diazinon 50W in 25 gals. of water and apply as an overall spray to ceilings and walls of livestock sheds, calf barns, hog barns, feeding sheds, and other farm buildings except poultry houses, dairy barns, and milk rooms. Areas where flies congregate, such as garbage dumps and corrals, should be sprayed thoroughly. Longer residual control may be obtained by increasing the rate to 4 lbs. in 25 gals. of water. Applications should be made to permit run-off. One gal. of the above spray should cover from 350-750 sq. ft. of area, depending on the surface treated. Repeat applications as fly populations warrant.

Bait Sprays: Mix the appropriate amount of Diazinon 50W with 1 lb. of sugar (or 2 cups of syrup or molasses) for the desired application.

For knapsack or similar type sprayer, mix 1/2 lb. with the sugar in 2 1/2 gals. of water and spray areas frequented by flies including cracks, crevices, doorways, windows, and other areas where flies congregate. Repeat applications as fly populations warrant.

For spraying car application, mix 2 oz. with the sugar in 5 gals. of water and spray lightly over the floor in areas frequented by flies. Applications every day or two are generally required.

Do not contaminate milk, food, drinking water, or apply to animals or poultry. Do not use in edible product areas of food processing plants or other areas where food is commercially prepared or processed. Do not use in dwellings.

NET CONTENTS

11/83

LOT NO.

Manufactured by



Drug & Chemical Co., Inc.

Plant: Kean Road, Sandersville, GA. 31082
 Office: C.B. 2000 Floral Park, N.Y. 11001

655-456