

655-459

11-17-2003

1/2



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

NOV 17 2003

Mr. Jeffrey Miller
Prentiss Incorporated
C.B. 2000
Floral Park, NY 11002-2000

Dear Mr. Miller:

Subject: Add California Supplemental label
Prentox Diazinon AG-500 Insecticide
EPA Registration Number 655-459
Your submission dated October 14, 2003

The labeling referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(a), is acceptable and a stamped copy of the California Supplemental label is enclosed for your records.

Sincerely,

George LaRocca.
Product Manager 13
Insecticide Branch
Registration Division (7505C)

SUPPLEMENTAL LABEL

RESTRICTED USE PESTICIDE
DUE TO AVIAN AND AQUATIC TOXICITY
FOR RETAIL SALE TO AND USE BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT
SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION

PRENTOX

DIAZINON AG-500 INSECTICIDE

EPA Reg. No. 655-459

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL AND PAMPHLET, USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

FOR PROTECTION OF AQUATIC SPECIES IN COUNTIES AND PORTIONS OF COUNTIES LOCATED IN THE SACRAMENTO AND SAN JOAQUIN VALLEYS BELOW 1000 FEET ELEVATION FROM POTENTIAL CONTAMINATION IN SURFACE WATERS OF DIAZINON APPLIED DURING THE DORMANT SPRAY PERIOD, THE FOLLOWING ADDITIONAL LABEL REQUIREMENTS APPLY:

- Dormant applications on orchard crops are restricted to ground application equipment only.
- Do not apply within 100 feet upslope of "sensitive aquatic sites" such as any irrigation ditch, drainage canal or body of water that may drain into a river or tributary unless a suitable method is used to contain or divert runoff waters. Waters that are contained or diverted must be held for a minimum of 72 hours before release into a sensitive aquatic site.
- Maintain a vegetative buffer strip a minimum of 10 feet wide from the edge of a field that is adjacent to and within 100 feet of sensitive aquatic sites.
- Do not apply this product to orchards when soil moisture is at field capacity, and/or when a storm event likely to produce runoff from the treated orchard is forecasted by NOAA/NWS, (National Weather Service) to occur within 48 hours following application.
- Make dormant applications only when insect scouting information or the recommendation of a Pest Control Advisor indicate treatment is required. (See UC IPM Guidelines for San Jose Scale in stone fruits. Use the prune dormant spur sampling program to determine need for a dormant treatment in that crop.)
- Apply only when wind speed is 3-10 mph at the application site as measured by an anemometer outside of the orchard on the side nearest and upwind from a sensitive site.
- When sensitive aquatic sites are downwind from orchards, spray the first three rows nearest the sensitive sites only when the wind is blowing away from the sites. The row at the edge of the field next to sensitive aquatic sites must be sprayed with the outside nozzles turned off. Spray must not be directed higher than the tree canopy and spray must be directed away from sensitive aquatic sites.
- The Stewardship Bulletin "Orchard Practices for Protecting Surface Water", must be available to handlers and equipment operators at the application site during all application activities. (note: Bulletin available through CURES)

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION. PLEASE REFER TO CONTAINER LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Manufactured by:

PRENTISS INCORPORATED

Office: C.B. 2000, Floral Park, NY 11002-2000
Plant: Kaolin Road, Sandersville, GA 31082

ACCEPTED

NOV 17 2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 655-459

2/2