

Meat, poultry, and other food processing plants, refuse areas and containers.

SPRAY RECOMMENDATIONS			
RONNEL %	AMOUNT RONNEL 24E FOR		
	1 GAL.	25 GAL.	100 GAL.
1%	5-1/3 oz.	1 gal.	4 gal.

Remarks: Do not apply in the edible product areas of food processing plants.

Restaurants, and supermarkets. (Outside area).

1/2% or 1%	2-2/3 oz. or 5-1/3 oz.	2 qt. or 1 gal.	2 gal. or 4 gal.
------------	------------------------	-----------------	------------------

Remarks: Spray outside walls, receiving docks, doorways and refuse areas.

Dwellings - Outside areas, including porches, patios and garages.

1/2% or 1%	2-2/3 oz. or 5-1/3 oz.	2 qt. or 1 gal.	2 gal. or 4 gal.
------------	------------------------	-----------------	------------------

Remarks: Spray to point of runoff.

Caged Layers - on wire.

2%	10-2/3 oz.	2 gal.	8 gal.
----	------------	--------	--------

Remarks: Spray poultry droppings beneath cages.

FLY BAIT - Where flies congregate: floors, windows and doorways.

2%	10-2/3 oz.	2 gal.	8 gal.
----	------------	--------	--------

Remarks: Add 3 oz. of sugar per gallon diluted spray. Spray or sprinkle at 1 to 2 day intervals until control is obtained. Do not use in food handling or food processing areas.

DARKLING BEETLES - in poultry houses.

1/2%	2-2/3 oz.	2 qt.	2 gal.
------	-----------	-------	--------

Remarks: Spray floors, footings, and lower 2 ft. of inside walls of poultry houses. Use 2 gals. of spray per 100 sq. ft. of area. Allow spray to dry before birds are placed on litter. Re-treat as necessary, but remove birds from area until spray is dry.

COCKROACH CONTROL - Use 1% spray for German roaches, 2% for others.

Remarks: Apply as a spot spray to all areas frequented by cockroaches.

Buildings - Dwellings, institutions and food handling establishments.

1% or 2%	5-1/3 oz. or 10-2/3 oz.	1 gal. or 2 gal.	4 gal. or 8 gal.
----------	-------------------------	------------------	------------------

Remarks: Apply as a spot spray to all areas frequented by cockroaches. Do not use in edible product areas, or in serving areas while food is exposed.

Food Processing Plants

1% or 2%	5-1/3 oz. or 10-2/3 oz.	1 gal. or 2 gal.	4 gals. or 8 gals.
----------	-------------------------	------------------	--------------------

Remarks: Do not apply in the edible product areas of food processing plants.

Meat Packing and Poultry Processing Plants.

SPRAY RECOMMENDATIONS			
RONNEL %	AMOUNT RONNEL 24E FOR		
	1 GAL.	25 GAL.	100 GAL.
2%	10-2/3 oz.	2 gal.	8 gal.

Remarks: Do not apply in the edible products areas of food processing plants.

MISCELLANEOUS INSECTS:

Fleas and spiders in buildings, laundry trucks, and dog kennels.

1%	5-1/3 oz.	1 gal.	4 gal.
----	-----------	--------	--------

Remarks: Apply 1 gallon of dilute spray per 1000 sq. ft. Spray floors, walls, cracks, crevices, sleeping quarters of pets, etc., where insects breed and harbor. Do not use in food handling or food processing areas.

Bedbugs in buildings and sleeping quarters, on beds and bedding.

1%	5-1/3 oz.	1 gal.	4 gal.
----	-----------	--------	--------

Remarks: Spray mattress and bedding lightly, including seams and folds, cracks and crevices in bed frames, walls, baseboards, and floors. Allow bedding and mattress to dry before using.

Fleas in lawns and kennel runs.

1%	5-1/3 oz.	1 gal.	4 gal.
----	-----------	--------	--------

Remarks: Spray infested areas. Keep children off treated area until spray has dried.

Ants, crickets, mosquitoes and silverfish.

1%	5-1/3 oz.	1 gal.	4 gal.
----	-----------	--------	--------

Remarks: Spray directly on insects where possible. Spray surfaces where insects alight, breed and congregate, such as under sinks and drainboards, in cabinets, and around baseboards, door sills and frames, window frames, garages, and outbuildings. Repeat as necessary.

Chinch Bugs in lawns (Florida only): Retreat as necessary. Thin Turf or Thick Turf (thatch and organic matter 1/2 inch thick or more).

0.15%	0.8 oz.	20 oz.	80 oz.
-------	---------	--------	--------

Remarks: Apply following a good rain or a drench with 1/2 to 1 inch of water. Dilute with water only. Use 25 gallons diluted spray per 1250 sq. ft. For thick turf, repeat in 2 weeks, if necessary.

NOTICE:

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with Directions under normal conditions of use, but neither this warranty nor any other warranty or merchantability or fitness for a particular purpose express or implied extends to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use.

A PRENTISS PEST-TESTED PRODUCT

Prentox

RONNEL 24E INSECTICIDE

FOR FLIES, LICE & PEST CONTROL USE

ACTIVE INGREDIENTS:

Ronnel - O,O-dimethyl O-(2,4,5-trichlorophenyl) phosphorothioate	24.0%
Aromatic Petroleum derivative solvent	58.9%

INERT INGREDIENTS:	17.1%
TOTAL:	100.0%

CAUTION
KEEP OUT OF REACH OF CHILDREN

READ COMPLETE PRECAUTIONS ON SIDE PANEL

FOR USE ON: Cattle, Hogs, Sheep, Goats, Newborn too.

FOR USE IN: Dairy Barns, Milk Rooms, Hog Barns, Poultry Houses, Farm Buildings, Backrubbers.

TO CONTROL: Lice, Horn Flies, Face Flies, Screw-worms, Wool Maggots, Sheep Keds, Cockroaches, House Flies, Stable Flies, Mosquitoes, Spiders, Ants, Ticks.

PRENTOX RONNEL 24E - Livestock remedy. Not for human use.

WATER EMULSIFIABLE - OIL SOLUBLE

TAKE TIME OBSERVE LABEL DIRECTIONS!

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME

1 GALLON

MANUFACTURED BY:

E.P.A. Reg. No. 655-472

Prentiss Drug & Chemical Co., Inc.

PLANT: 338 WILSON AVE. NEWARK, N.J. 07105 • OFFICE: 163 JEROME AVE. NEW YORK, N.Y. 10017

CAUTION
KEEP OUT OF REACH OF CHILDREN

INJURIOUS TO THE EYES. IRRITATING TO THE SKIN. HARMFUL IF SWALLOWED. COMBUSTIBLE.

Do not get in eyes. Avoid contact with the skin. Avoid prolonged breathing of spray mists. Avoid contamination of food, feedstuffs and water supplies. Keep material out of reach of animals. Keep away from heat and open flame. Keep container closed. Wash thoroughly with soap and water after handling.

NOTE TO PHYSICIAN:

This material is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

USE PRECAUTIONS:

Do not spray sheep and goats within 28 days of slaughter. Do not spray beef cattle with 3/4% Ronnel within 14 days of slaughter. Do not spray beef cattle with 1/2% Ronnel within 7 days of slaughter. Beef cattle may be sprayed with 1/4% Ronnel any time prior to slaughter. Backrubber may be used any time on beef or dairy cattle. Do not use spray on animals producing milk for human consumption. Dairy cows should be treated early in their dry period. Do not spray dairy animals with 3/4% Ronnel for 14 days prior to freshening. If dairy cows or heifers freshen within 14 days of the last spray treatment, milk must not be used for food for the required 14 days after the last spray treatment with 3/4% Ronnel. Do not spray dairy animals with 1/4% or 1/2% Ronnel for 7 days prior to freshening. If dairy cows or heifers freshen within 7 days of last treatment, milk must not be used for food for the required 7 days after the last spray treatment with 1/4% or 1/2% Ronnel. Remove lactating dairy animals from buildings while applying residual fly sprays. Do not allow insecticide to come in contact with food, feed or water supplies.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY PERFORATING OR CRUSHING IT. BURY OR DISCARD IT AT SAFE PLACE, AWAY FROM WATER SUPPLIES.

WARNING

Ronnel is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with, or exposure to cholinesterase inhibiting drugs, pesticides or chemicals. Treatment with excessive dosages of organo-phosphates may lead to the general symptoms of muscular weakness, diarrhea and excessive salivation. At the first sign of adverse reaction, consult a veterinarian. Atropine is an antidote.

CAUTION
COMBUSTIBLE MIXTURE

BACKRUBBER RECOMMENDATIONS

Cattle Backrubber — For hornflies. Aids in the control of lice if used continuously. Aids in the control of face flies if backrubber is placed so animals can rub face. Place where animals regularly travel or congregate.

Mix 1 gal. RONNEL 24E with 27 gal. diesel or mineral oil. Saturate fibre. Recharge as necessary. Do not mix with used motor oil.

PROBLEM — LIVESTOCK INSECTS:

Cattle and Calves: Hornflies, ear ticks and screw-worms.

SPRAY RECOMMENDATIONS

RONNEL %	AMOUNT RONNEL 24E FOR		
	1 GAL.	25 GAL.	100 GAL.
1/2%	2-2/3 oz.	2 qts.	2 gals.

Lice

1/4%	1-1/3 oz.	1 qt.	1 gal.
------	-----------	-------	--------

Lone Star & Winter Ticks

3/4%	4 oz.	3 qt.	3 gal.
------	-------	-------	--------

Sheep, Goats, Lambs, Kids: Hornflies, screw-worms and (wool maggots), fleeceworms.

1/2%	2-2/3 oz.	2 qt.	2 gal.
------	-----------	-------	--------

Lice and sheep keds.

1/4%	1-1/3 oz.	1 qt.	1 gal.
------	-----------	-------	--------

Hogs: Lice

1/4%	1-1/3 oz.	1 qt.	1 gal.
------	-----------	-------	--------

Remarks: Dilute with water only. Spray entire animal to wet thoroughly. Retreat as necessary, but not more than every 2 weeks.

For ear ticks, spray ears and surrounding area thoroughly.

RESIDUAL FLY SPRAY — A 1/2% Spray will give about two weeks control. A 1% Spray will give 4-6 weeks control.

Remarks: Spray walls, ceilings, partitions to obtain complete coverage with no runoff. Use 1 gal. of diluted spray per 500-1000 sq. ft.

Farm Buildings — Poultry houses, livestock shelters, milk rooms, dairy and hog barns.

SPRAY RECOMMENDATIONS

RONNEL %	AMOUNT RONNEL 24E FOR		
	1 GAL.	25 GAL.	100 GAL.
1/2% or 1%	2-2/3 oz. or 5-1/3 oz.	2 qts. or 1 gal.	2 gals. or 4 gals.

Remarks: Add 3 oz. of sugar per gallon for added effectiveness. To prevent contamination keep milk and milking equipment covered while spraying. Do not spray directly on birds.