

1973

GENERAL INSECTICIDE
 FEDERAL INSECTICIDE ACT
 DIAZINON REG. STER.
 655-475

| FOR APPLYING | |
|--|---|
| RATE & AREA COVERED | NOTES |
| 2 tbsps. (1 fl. oz.) in 3 gals. of water covers 125 sq. ft. 16 fl. oz. in 24 gal. of water covers 1,000 sq. ft. | Thoroughly wet down grass a few hours before applying. For Lawn Billbugs, water lawn after treatment. |
| 4 tbsps. (2 fl. oz.) per 3 gals. water covers 125 sq. ft. 32 oz./24 gals. of water covers 1,000 sq. ft. | In the north, if CHINCHBUGS are the only insect problem, a lower rate of 2 tbsps. in 3 gals. of water covers 250 sq. ft. |
| 2 tbsps. (1 fl. oz.) per 3 gals. of water. Apply as a thorough foliar spray. | Spray underside of leaves. Wet plants to the dripping point. |
| 2 tbsps. (1 fl. oz.) per 3 gals. of water. Apply as a thorough foliar spray. | Wet leaves and branches to the dripping point. Try to penetrate dense foliage. |
| 2 tbsps. (1 fl. oz.) per 3 gals. of water. Apply as a full cover spray. | Start spraying after blooming and repeat at 10 to 14 day intervals if needed. Do not spray fruit within 21 days of picking. |
| 2 tbsps. (1 fl. oz.) per 3 gals. of water covers 1,000 sq. ft. | Generally applications at 7-10 day intervals may be required. Do not apply within 1 day of picking tomatoes or 14 days of picking all other crops listed in this section. |
| Apply 1½ fl. oz. per 100 sq. ft. as a soil treatment in sufficient water to obtain thorough coverage. | Make application to soil surface as a broadcast spray prior to planting. Thoroughly mix the chemical into the top 4-8 inches of soil. |
| Add 8 fl. oz. per 3 gals. of water. | Thoroughly spray window frames, doors, sills, screens, porch and patio walls and ceiling, foundations, walks, garbage cans and into cracks. Do not treat plants or animals with this dosage rate. |

HOW TO MIX AND APPLY

NUMBER ONE:


Determine size of spraying job. For lawns and vegetables, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear, or in advance of an expected infestation. Repeat applications as needed or as noted in the rate chart.

NUMBER TWO:

Use clean sprayer. Carefully measure and mix the amount of Diazinon and water as suggested in the chart.

NUMBER THREE:

Spray as directed in the rate chart. Use coarse drenching sprays for lawns, soil and other surface applications. Use medium fine mist sprays for trees and plants. Thorough coverage is important. Flush sprayer with clean water after each use.



DIAZINON®*

25% EMULSIFIABLE CONCENTRATE

Lawn & Garden Insect Control

ACTIVE INGREDIENTS:

O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate* 25.0%
 Aromatic Petroleum Derivative Solvent 55.7%

INERT INGREDIENTS: 19.3%

TOTAL: 100.0%

DIAZINON® is a registered trademark of CIBA-GEIGY CORPORATION.

WARNING
 Keep out of reach of children

THIS PRODUCT IS HIGHLY TOXIC TO BEES EXPOSED TO DIRECT TREATMENT OR RESIDUES ON CROPS. PROTECTIVE INFORMATION MAY BE OBTAINED FROM YOUR COOPERATIVE AGRICULTURAL EXTENSION SERVICE.

SEE SIDE PANEL FOR ADDITIONAL WARNINGS.

KILLS CERTAIN outdoor insects which destroy lawns, roses, flowers, trees and shrubs or annoy pets and people. Contains Diazinon Insecticide which provides positive control of many "hard to kill" sucking and chewing insects.

E.P.A. Reg. No. 655-475

NET CONTENTS _____

Prentiss Drug & Chemical Co., Inc.
 PLANT 338 WILSON AVE NEWARK, N. J. 07105 • OFFICE 363 SEVENTH AVE NEW YORK, N. Y. 10001



DIAZINON®*

25% EMULSIFIABLE CONCENTRATE Lawn & Garden Insect Control

ACTIVE INGREDIENTS:

O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate* 25.0%
Aromatic Petroleum Derivative Solvent 55.7%

INERT INGREDIENTS: 19.3%

TOTAL: 100.0%

DIAZINON® is a registered trademark of CIBA-GEIGY CORPORATION.

WARNING
Keep out of reach of children

THIS PRODUCT IS HIGHLY TOXIC TO BEES EXPOSED TO DIRECT TREATMENT OR RESIDUES ON CROPS. PROTECTIVE INFORMATION MAY BE OBTAINED FROM YOUR COOPERATIVE AGRICULTURAL EXTENSION SERVICE.

SEE SIDE PANEL FOR ADDITIONAL WARNINGS.

KILLS CERTAIN outdoor insects which destroy lawns, roses, flowers, trees and shrubs or annoy pets and people. Contains Diazinon Insecticide which provides positive control of many "hard to kill" sucking and chewing insects.

E.P.A. Reg. No. 655-475

NET CONTENTS _____

Prentiss Drug & Chemical Co., Inc.

PLANT 338 WILSON AVE NEWARK, N. J. 07105 • OFFICE 363 SEVENTH AVE NEW YORK, N. Y. 10018

DO NOT USE IN THE HOME

WARNING:

Keep out of reach of children. May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid contamination of feed and food-stuffs.

Do not use on household pets or humans. Do not permit children or pets to go on to the sprayed grass until the spray has completely dried. Keep out of reach of domestic animals.

DO NOT STORE OR USE NEAR HEAT OR OPEN FLAME.

DONOT CONTAMINATE ORNAMENTAL FISHPONDS.

FISH AND WILDLIFE WARNING:

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated area. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

WARNING:

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NOTICE:

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.



RATE CHART

| PLANT | INSECT PESTS |
|--|--|
| LAWNS: Bromegrass, Bluegrass, St. Augustine, Dichandra, Bermuda | Aphids, Mites, Leafhoppers, Thrips |
| ROSES & FLOWERS: Roses, Azaleas, Chrysanthemums | Aphids, Mites, Scale Crawlers, Leaf Miners, Bagworms |
| TREES & SHRUBS: Oak, Maple, Locust, Holly, Boxwood, Pine | Codling Moths, Mites, Aphids, Leaf Rollers, Scale Crawlers |
| FRUITS: Apples, Peaches, Oranges, Plums, Grapes, Cherries, Strawberries | Bean Beetles, Aphids, Leaf Miners, Cabbage-worms, Cucumber Beetles |
| VEGETABLES: Tomatoes, Carrots, Beans, Onions, Cucumbers, Cantaloupes, Peppers, Cabbage | Cutworms (Surface and Subterranean) Wireworms, Root Maggots, Mole Crickets |
| Cabbage, Carrots, Corn, Lettuce, Peas, Radishes, Red Beets, Snap Beans, Tomatoes, Turnips | Flies, Ants, Cockroaches, Boxelder bugs, Clover Mites, Spiders, Ticks, Chiggers, Earwigs |
| Nuisance pests in outside areas | |

DO NOT USE ON CERTAIN FERNS, HIBISCUS, POINSETTIA, PAPAYA, PILEA AND GARDENIA AND ORCHIDS, SINCE INJURY MAY OCCUR.