

Prentox PCO Pyrethrum Powder 0.9%
EPA Registration No. 655-479

PRENTOX® PCO PYRETHRUM POWDER 0.9%

For the control of Cockroaches, Ants & Silverfish — in Homes, Non-Food areas of Restaurants, and Food Processing Plants. Also for the control of Lice on Poultry and Pet Birds, Fleas and Ticks on Premises, Fleas and Lice on Dogs, Cats, Fleas on Gerbils, Hamsters and White Mice.

ACTIVE INGREDIENTS:

Pyrethrins	0.9%
INERT OTHER INGREDIENTS*:	99.1%
Total:	100.0%

*(Powdered Pyrethrum Flowers other than Pyrethrins)
PRENTOX® - Registered Trademark of Prentiss Incorporated

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT FIRST AID

If in eyes — Flush immediately with plenty of water. Get medical attention if irritation persists.
If on skin — Wash thoroughly with soap and water.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 655-479

EPA EST. NO. 655-GA-1

NET CONTENTS _____

LOT NO. _____

Manufactured by:

PRENTISS INCORPORATED

Plant: Kaolin Road, Sandersville, GA 31082
Office: C.B. 2000, Floral Park, NY 11002-2000

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Avoid breathing dust. Avoid contact with eyes. In case of contact, immediately flush eyes with plenty of water. Get medical attention if irritation persists. Wash hands thoroughly after use. This product contains dried, finely ground flowers of the species *chrysanthemum cinerariaefolium*, and inhalation exposure may cause allergic reactions similar to hay fever. May also aggravate conditions such as asthma.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE: Do not apply where children are likely to come in frequent contact with treated surfaces. Do not contaminate food, food containers or cooking utensils. Repeat application as necessary.

Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

FOR CONTROL OF ROACHES: Dust thoroughly all areas where roaches crawl and hide, around water pipes, in cracks and crevices, under and behind cabinets, refrigerators and sinks, and around doors, windows and bookcases. Repeat application as necessary.

FOR CONTROL OF SILVERFISH: Dust thoroughly in cracks and crevices around baseboards, door frames, bookcases, and storage areas.

FOR CONTROL OF ANTS: Dust thoroughly around door and window frames, ant trails, and ant hills.

LICE ON POULTRY & PET BIRDS: Dust the powder carefully into the feathers, under the wings, along the back and around the vent. Rub into feathers of head. Nests, roosts, and runways, porches and cages should be thoroughly dusted. Destroy infested food or cuttlebone. Repeat as needed.

FLEAS & TICKS ON PREMISES: Dust thoroughly all infested areas, particularly crevices in woodwork, floor and floor covering, and animal sleeping quarters. Repeat as needed.

ACCEPTED

MAY 26 1998

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 655-479

PETS

PRE-TREATMENT DIRECTIONS

READ ENTIRE LABEL BEFORE EACH USE

USE ONLY ON DOGS TO KILL FLEAS, LICE AND BROWN DOG TICKS: Do not use on dogs under 12 weeks.

SPECIAL ANIMALS

Consult a veterinarian or call the National Pesticide Telecommunications Network at 1-800-858-7378, before using this product on debilitated, aged, pregnant, nursing or medicated animals.

ADVERSE REACTION INFORMATION

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your dog with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately, or call the National Pesticide Telecommunications Network at 1-800-858-7378.

APPLICATION DIRECTIONS

FLEAS, LICE AND BROWN DOG TICKS ON CATS & DOGS: Dust the powder onto the animal's hair, including head (except eyes, mouth and nose), back, sides, legs, ears, and tail and work it down to the skin. Avoid any sores. After treatment, brush or comb the animal's coat to remove dead fleas and lice. Repeat as needed. Dust sleeping quarters and replace bedding if necessary.

REAPPLICATION DIRECTIONS

Do not reapply product for 7 days.

BEDBUGS: Dismantle beds completely and dust all parts thoroughly, including mattress seams and box springs. Dust all baseboards and wall cracks thoroughly. Repeat as needed.

FLEAS ON GERBILS, HAMSTERS AND WHITE MICE: Do not use if female is pregnant or nursing young. Dust the powder onto the animal's hair and work it down to the skin. After treatment, brush or comb the animal's coat to remove dead fleas and lice. Repeat as needed.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

STORAGE: If bag is damaged, stop the spill by repositioning the bag or otherwise repairing it. Take care to avoid contact with pesticide and wear protective gear. Do not breathe dust. Sweep spill on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities by burning. If burned, stay out of smoke.